

GEOMETRY OF NILMANIFOLDS WITH LEFT-INVARIANT COMPLEX STRUCTURE AND DEFORMATIONS IN THE LARGE

SÖNKE ROLLENSKE

ABSTRACT. The relation between nilmanifolds with left-invariant complex structure and iterated principal holomorphic torus bundles is clarified and we give criteria under which deformations in the large are again of such type. As an application we obtain a fairly complete picture in complex dimension three.

AMS Subject classification: 32G05; (32G08, 17B30, 53C30)

CONTENTS

Introduction	1
1. Nilpotent Lie-algebras and nilmanifolds with left-invariant complex structure	5
1.1. Lie-algebras with a complex structure	5
1.2. Nilmanifolds with left-invariant complex structure	6
1.3. Dolbeault cohomology of nilmanifolds and small deformations	11
1.4. Examples and Counterexamples	12
2. Albanese-Quotients and deformations in the large	15
2.1. Definitions and results	15
2.2. Proof of Theorem 2.11	19
3. Complex structures on certain Lie-algebras	25
3.1. Notations and basic results	25
3.2. The case $\dim(\mathcal{C}^1\mathfrak{g}) = 1$	27
3.3. The case $\dim(\mathcal{C}^1\mathfrak{g}) = 2$	29
3.4. The case $\dim(\mathcal{C}^1\mathfrak{g}) = 3$	30
4. Applications	35
4.1. The Main Theorem	35
4.2. Deformations and geometric structure in dimension 3	37
References	39

INTRODUCTION

A very general question in the theory of complex manifolds is the following: let M be a compact, differentiable manifold and let

$$\mathcal{C} := \{J \in \text{End}(TM) \mid J^2 = -id_{TM}, J \text{ a complex structure}\}$$

be the space of all complex structures on M . What can we say about $\mathcal{C}$ and its connected components?

If $\mathcal{C}$ is non-empty consider a compact, complex manifold $X = (M, J)$. The theory developed by Kodaira and Spencer in the 50's [KS58] and culminating in the theorem of Kuranishi [Kur62] succeeds in giving a rather precise description of a slice of $\mathcal{C}$ containing J which is transversal to the orbit of the natural action of $\text{Diff}^+(M)$, called the Kuranishi slice.

While we have this powerful tool for the study of small deformations there is no general method available to study the connected components of $\mathcal{C}$.

From another point of view we say that two compact, complex manifolds X and X' are directly deformation equivalent $X \sim_{\text{def}} X'$ if there exists an irreducible, flat family $\pi : \mathcal{M} \rightarrow \mathcal{B}$ of compact, complex manifolds over a complex analytic space $\mathcal{B}$ such that $X \cong \pi^{-1}(b)$ and $X' \cong \pi^{-1}(b')$ for some points $b, b' \in \mathcal{B}$. The manifold X is said to be a deformation in the large of X' if both are in the same equivalence class with respect to the equivalence relation generated by $\sim_{\text{def}}$, which is the case if and only if both are in the same connected component of $\mathcal{C}$.

Even the seemingly natural fact that any deformation in the large of a complex torus is again a complex torus has been fully proved only in 2002 by Catanese [Cat02]. In [Cat04] he studies more in general deformations in the large of principal holomorphic torus bundles, especially bundles of elliptic curves. This was the starting point for our research.

It turns out (see [CF06]) that the right context to generalise Catanese's results is the theory of left-invariant complex structures on nilmanifolds, i.e., compact quotients of nilpotent real Lie-groups by discrete subgroups.

Nilmanifolds with left-invariant complex structure provide an important source for examples in complex differential geometry. Among these are the so-called Kodaira-Thurston manifolds, historically the first examples known to admit both a complex structure and a symplectic structure but no Kähler structure. In fact, a nilmanifold M admits a Kähler structure if and only if it is a complex torus [BG88, Has89] and the author showed [Rol07c] that nilmanifolds can be arbitrarily far from Kähler manifolds in the sense that the Frölicher spectral sequence may be arbitrarily non-degenerate.

But unfortunately, even if every (iterated) principal holomorphic torus bundle can be regarded as a nilmanifold, the converse is far from true. Moreover it turns out that even a small deformation of a principal holomorphic torus bundle may not admit such a structure (see Example 1.14). Small deformations of nilmanifolds were already studied for so-called abelian complex structures in [CFP06, MPPS06] and for general left-invariant complex structure on nilmanifolds in [Rol08b]. Thus we are concerned with these two problems:

- Give conditions under which a left-invariant complex structure on a nilmanifold gives rise to a structure of iterated (principal) holomorphic torus bundle.
- Study deformations in the large of iterated (principal) holomorphic torus bundles.

There is already a vast literature concerning nilpotent Lie-algebras and left-invariant complex structures on nilmanifolds (see e.g. the articles of Console, Cordero, Fernandez, Fino, Grantcharov, Gray, McLaughlin, Pedersen, Poon, Salamon, Ugarte, et al. cited in the bibliography) and we

recapitulate the basic results in Section 1, emphasising the complex geometric structure. In Section 1.3 we will also recall the results on Dolbeault-cohomology and small deformations.

A nilmanifold M can be characterised by a triple $(\mathfrak{g}, J, \Gamma)$ where $\mathfrak{g}$ is the nilpotent Lie-algebra associated to a simply connected nilpotent Lie-group G , J is an integrable complex structure on $\mathfrak{g}$ and $\Gamma \subset G$ is a (cocompact) lattice. The datum of either $\mathfrak{g}$ or Γ (considered as an abstract group) determines G up to unique isomorphism. The general philosophy is that the geometry of the compact, complex manifold $M_J = (\Gamma \backslash G, J)$ should be completely determined by the linear algebra of $\mathfrak{g}$, J and the $\mathbb{Q}$ -subalgebra generated by $\log \Gamma \subset \mathfrak{g}$. By abuse of notation we will sometimes write $M_J = (\mathfrak{g}, J, \Gamma)$.

The real geometry of nilmanifolds is well understood but only the existence of what we call a stable torus bundle series in the Lie-algebra $\mathfrak{g}$ (Definition 1.8) gives us sufficient control over the complex geometry of (M, J) . Geometrically this notion means the following:

On any real nilpotent Lie-group G there is a filtration by normal subgroups, e.g., the ascending central series,

$$G \supset H_s \supset \cdots \supset H_1 \supset H_0 = \{1\}$$

such that H_k/H_{k-1} is abelian and $\Gamma \cap H_k$ is a lattice in H_k for any lattice $\Gamma \subset G$.

In other words, any compact nilmanifold $M = \Gamma \backslash G$ can be represented as a tower of differentiable torus bundles $\pi_k : M_k \rightarrow M_{k+1}$

$$\begin{array}{ccc} T_1 & \hookrightarrow & M_1 = M \\ & & \downarrow \pi_1 \\ T_2 & \hookrightarrow & M_2 \\ & & \downarrow \pi_2 \\ & & \vdots \\ T_{s-1} & \hookrightarrow & M_{s-1} \\ & & \downarrow \pi_{s-1} \\ & & M_s = T_s \end{array}$$

where the compact torus T_k is the quotient of H_k/H_{k-1} by the lattice and M_k is the compact nilmanifold obtained from the Lie-group G/H_{k-1} by taking the quotient with respect to the image of Γ .

Such a decomposition into an iterated principal bundle is far from unique. If the subgroups can be chosen in such a way that for every left-invariant complex structure J on M we have

- (i) The complex structure J induces a left-invariant complex structure on M_k for all k .
- (ii) All the maps $\pi_k : M_k \rightarrow M_{k+1}$ are holomorphic with respect to these complex structures.

then we say that $\mathfrak{g}$ admits a stable torus bundle series.

This phenomenon is not as uncommon as it may seem at first sight; for example every principal holomorphic torus bundle over an elliptic curve which is not a product has this property (Example 1.13). Section 3 will be devoted to giving a fairly complete picture of the occurring cases if $\dim[\mathfrak{g}, \mathfrak{g}] \leq 3$. A large part of the classification of complex structures on real 6-dimension nilpotent Lie-algebras [Sal01, Uga04] can be recovered from our more general results.

It turns out that finding a stable torus bundle series (STBS) for some nilmanifold M_J is a good step on the way to prove that every deformation in the large of M_J is again such a nilmanifold. Indeed, in this case the holomorphic fibration over a torus $\pi : M \rightarrow T_s$ can be realised as a (topologically) fixed quotient of the Albanese variety and this will enable us in Section 2 to determine all deformations in the large if the fibres of π have sufficiently nice properties:

Theorem 2.12 — *Let G be a simply connected nilpotent Lie-group with Lie-algebra $\mathfrak{g}$ and let $\Gamma \subset G$ be a lattice such that the following holds:*

- (i) $\mathfrak{g}$ admits a stable torus bundle series $(\mathcal{S}^i \mathfrak{g})_{i=0, \dots, t}$ (cf. Definition 1.8).
- (ii) Nilmanifolds with Lie algebra $\mathcal{S}^{t-1} \mathfrak{g}$ and fundamental group $\Gamma \cap \exp(\mathcal{S}^{t-1} \mathfrak{g})$ constitute a good fibre class (cf. Definition 2.9).

Then any deformation in the large of a nilmanifold with left-invariant complex structure $M = (\Gamma \backslash G, J)$ is again of the same type.

Generalising the methods used in [Cat04] we will have to deal with the fact that the dimension of the Albanese variety may vary in a family of nilmanifolds.

In Section 4 we apply our results on the deformation of complex structures to the classification obtained in Section 3 and thus give several series of examples (in arbitrary dimension) for which left-invariant complex structures yield a union of connected components of the space of complex structures $\mathcal{C}$. This holds for example if $\dim[\mathfrak{g}, \mathfrak{g}] \leq 1$ or if $\mathfrak{g}$ is 3-step nilpotent and $\dim[\mathfrak{g}, \mathfrak{g}] = 2$. We refer to Theorem 4.1 and Theorem 4.3 for a complete list of cases.

There are only finitely many real 6-dimensional, nilpotent Lie-algebras that admit a complex structure [Mag86, Sal01] and we obtain a quite complete picture of their geometry and deformations in Section 4.2:

Theorem 4.4 — *Let $M = (\mathfrak{g}, J, \Gamma)$ be a complex 3-dimensional nilmanifold with left-invariant complex structure.*

- (i) *If $\mathfrak{g}$ is not in $\{\mathfrak{h}_7, \mathfrak{h}_{19}^-, \mathfrak{h}_{26}^+\}$, then $\mathfrak{g}$ admits a SPTBS and hence M_J has the structure of an iterated principal holomorphic torus bundle. We list the possibilities in the following table:*

base	fibre	corresponding Lie-algebras
3-torus	-	$\mathfrak{h}_1$
2-torus	elliptic curve	$\mathfrak{h}_2, \mathfrak{h}_3, \mathfrak{h}_4, \mathfrak{h}_5, \mathfrak{h}_6$
elliptic curve	2-torus	$\mathfrak{h}_8$
Kodaira surface	elliptic curve	$\mathfrak{h}_9, \mathfrak{h}_{10}, \mathfrak{h}_{11}, \mathfrak{h}_{12}, \mathfrak{h}_{13}, \mathfrak{h}_{14}, \mathfrak{h}_{15}, \mathfrak{h}_{16}$

Every deformation in the large is of the same type.

- (ii) *If $\mathfrak{g} = \mathfrak{h}_{19}^-$ or $\mathfrak{g} = \mathfrak{h}_{26}^+$ then $\mathfrak{g}$ admits a STBS and M_J can be described as 2-torus bundle over an elliptic curve but there is no principal torus bundle structure. Every deformation in the large is of the same type.*
- (iii) *If $\mathfrak{g} = \mathfrak{h}_7$ then there is a dense subset of the space of all left-invariant complex structures for which M admits the structure of principal holomorphic bundle of elliptic curves over a Kodaira surface but this is not true for all complex structures.*

It would also be interesting to study the space of left-invariant complex structures more in detail, for example determine when it is smooth or universal, and perhaps look for some kind of moduli space or a description of a connected component of the Teichmüller space. These questions have already been addressed in several cases by various authors [Cat04, CF06, GMPP04, KS04].

Nevertheless the conditions given in Theorem 2.11 are rather strong and we show in Example 1.14 that they do not need to be satisfied. This leads to the question: Which is the simplest example of a nilmanifold M_J with left-invariant complex structure such that not every deformation in the large carries a left-invariant complex structure?

Our theory of Albanese-Quotients in Section 2 is not restricted to nilmanifolds and it would be nice to find other applications.

Acknowledgements. This work is part of my PhD-thesis [Rol07a]. I would like to express my gratitude to my adviser Fabrizio Catanese for suggesting this research, constant encouragement and several useful and critical remarks. Simon Salamon gave valuable references to the literature and helped to improve the presentation at several points. Jörg Winkelmann remarked that every good Albanese variety is in fact very good. An invaluable bibliographic hint due to Oliver Goertsches opened a new perspective on the problem. The referee and the editor made several suggestion which helped to improve the presentation of the results.

I am grateful for the hospitality of the Max Planck Institut in Bonn where Section 2 got the final polishing; during the revision of the article I was visiting Imperial College London supported by a DFG Forschungsstipendium.

1. NILPOTENT LIE-ALGEBRAS AND NILMANIFOLDS WITH LEFT-INVARIANT COMPLEX STRUCTURE

In this section we will introduce the objects of our study and describe their basic properties. We will emphasise the complex-geometric structure of the nilmanifolds but the expert will find nothing new.

The geometrically important notion of stable (principal) holomorphic torus bundle series (SPTBS) will be given in Definition 1.8.

1.1. Lie-algebras with a complex structure. We will throughout need the yoga of almost complex structures and will now recall some basic definitions and notations.

Let $\mathfrak{g}$ be a finite dimensional real Lie-algebra and J an almost complex structure on the underlying real vector space, i.e., J is an endomorphism

of $\mathfrak{g}$ such that $J^2 = -Id_{\mathfrak{g}}$; setting $ix = Jx$ for $x \in \mathfrak{g}$ this makes $\mathfrak{g}$ into a complex vector space. We can decompose the complexified Lie-algebra $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g}^{1,0} \oplus \mathfrak{g}^{0,1}$ into the $\pm i$ -eigenspaces of the $\mathbb{C}$ -linear extension of J and every decomposition $\mathfrak{g}_{\mathbb{C}} = U \oplus \bar{U}$ gives rise to a unique almost complex structure J such that $\mathfrak{g}^{1,0} = U$.

Usually we will use small letters $x, y, \dots$ for elements of $\mathfrak{g}$ and capital letters $X, Y, \dots$ for elements in $\mathfrak{g}^{1,0}$; elements in $\mathfrak{g}^{0,1}$ will be denoted by $\bar{X}, \bar{Y}, \dots$ using complex conjugation.

The exterior algebra of the dual vector space $\mathfrak{g}^*$ decomposes as

$$\Lambda^k \mathfrak{g}_{\mathbb{C}}^* = \bigoplus_{p+q=k} \Lambda^p \mathfrak{g}^{*1,0} \otimes \Lambda^q \mathfrak{g}^{*0,1} = \bigoplus_{p+q=k} \Lambda^{p,q} \mathfrak{g}^*$$

and we have $\overline{\Lambda^{p,q} \mathfrak{g}^*} = \Lambda^{q,p} \mathfrak{g}^*$. A general reference for the linear algebra coming with a complex structure is [Huy05] (Section 1.2).

Definition 1.1 — *An almost complex structure J on a real Lie-algebra $\mathfrak{g}$ is said to be **integrable** if the Nijenhuis condition*

$$(1) \quad [x, y] - [Jx, Jy] + J[Jx, y] + J[x, Jy] = 0$$

holds for all $x, y \in \mathfrak{g}$ and in this case we call the pair $(\mathfrak{g}, J)$ a Lie-algebra with complex structure.

Hence by a complex structure on a Lie-algebra we will always mean an integrable one; otherwise we will speak of almost complex structures. It is easy to show that J is integrable if and only if $\mathfrak{g}^{1,0}$ is a subalgebra of $\mathfrak{g}_{\mathbb{C}}$ with the induced bracket.

If G is a real Lie-group with Lie-algebra $\mathfrak{g}$ then giving a left-invariant almost complex structure on G is equivalent to giving an almost complex structure J on $\mathfrak{g}$ and J is integrable if and only if it is integrable as an almost complex structure on G . It then induces a complex structure on G by the Newlander-Nirenberg theorem ([KN69], p.145) and G becomes a complex manifold. The elements of G act holomorphically by multiplication on the left but G is not a complex Lie-group in general.

1.2. Nilmanifolds with left-invariant complex structure. In this section we will collect a bunch of results on nilmanifolds, most of them well know and for which we claim no originality; but our presentation will emphasise the geometric structure of compact nilmanifolds.

If not otherwise stated $(\mathfrak{g}, J)$ will always be a Lie-algebra with (integrable) complex structure and G will be a associated simply connected Lie-group. By a torus we will always mean a compact torus.

Definition 1.2 — *A compact complex manifold $M_J := (M, J)$ is called **nil-manifold with left-invariant complex structure** if there is a nilpotent Lie algebra with complex structure $(\mathfrak{g}, J)$ and a lattice Γ in an associated simply-connected Lie-group such that $M_J \cong (\Gamma \backslash G, J)$.*

Since $\Gamma = \pi_1(M_J)$ determines G and hence $\mathfrak{g}$ up to isomorphism ([VGS00], p.45, Corollary 2.8), we will always identify M_J with $(\Gamma \backslash G, J)$ and call it a nilmanifolds with left-invariant complex structure of type $(\mathfrak{g}, \Gamma)$. By abuse of notation we will sometimes write $M_J = (\mathfrak{g}, J, \Gamma)$.

For nilpotent Lie-groups the exponential map $\exp : \mathfrak{g} \rightarrow G$ is a diffeomorphism and all analytic subgroups are closed and simply connected as well ([Var84], Theorem. 3.6.2, p. 196). The following often gives the possibility to use inductive arguments.

Lemma 1.3 — *Let $(\mathfrak{g}, J)$ be a nilpotent Lie-algebra with complex structure. Let $\mathfrak{h}$ be an ideal in $\mathfrak{g}$ such that $J\mathfrak{h} = \mathfrak{h}$. Let G and H be the associated simply connected Lie-groups endowed with the left-invariant complex structures induced by J . Then there is a holomorphic fibration $\pi : G \rightarrow G/H$ with fibre H .*

Proof. The map $\pi : G \rightarrow G/H$ is a real analytic fibration by the theory of Lie-groups and Lie-algebras since H is closed. Hence it remains to show that the differential of π is $\mathbb{C}$ -linear and since the complex structure is left-invariant it suffices to do so at the identity. But in this point the differential is given by the quotient map $\mathfrak{g} \rightarrow \mathfrak{g}/\mathfrak{h}$ which is $\mathbb{C}$ -linear by assumption. $\square$

Remark that we used the nilpotency of $\mathfrak{g}$ only to ensure that H is a closed subgroup.

1.2.1. *The real structure of $\Gamma \backslash G$.* We leave aside the complex structure for a moment and describe the geometry of the underlying real manifold. In the Lie-algebra $\mathfrak{g}$ we have the following filtrations:

- The descending central series (nilpotent series) is given by

$$C^0 \mathfrak{g} := \mathfrak{g}, \quad C^{i+1} \mathfrak{g} := [C^i \mathfrak{g}, \mathfrak{g}]$$

- The ascending central series is given by

$$Z^0 \mathfrak{g} := 0, \quad Z^{i+1} \mathfrak{g} := \{x \in \mathfrak{g} \mid [x, \mathfrak{g}] \subset Z^i \mathfrak{g}\}.$$

In particular $Z^1 \mathfrak{g} = Z \mathfrak{g}$ is the centre of $\mathfrak{g}$.

The Lie-algebra $\mathfrak{g}$ is called *s-step nilpotent* if $C^s \mathfrak{g} = 0$ and $C^{s-1} \mathfrak{g} \neq 0$ or equivalently $Z^s \mathfrak{g} = \mathfrak{g}$ and $Z^{s-1} \mathfrak{g} \subsetneq \mathfrak{g}$.

Let G_i be the simply connected nilpotent Lie-group associated to the Lie-algebra $\mathfrak{g}/Z^{i-1} \mathfrak{g}$ and let $F_i \cong \mathbb{R}^{n_i}$ be the abelian Lie-group corresponding to $Z^i \mathfrak{g}/Z^{i-1} \mathfrak{g}$. Then the map $G_i \rightarrow G_{i+1}$ is a principal bundle with fibre F_i and, if $\mathfrak{g}$ is *s-step nilpotent*, we get a tower of such bundles:

$$\begin{array}{ccc}
 F_1 \hookrightarrow & G_1 = G & \\
 & \downarrow & \\
 F_2 \hookrightarrow & G_2 & \\
 & \downarrow & \\
 & \vdots & \\
 & \downarrow & \\
 F_{s-1} \hookrightarrow & G_{s-1} & \\
 & \downarrow & \\
 & G_s = F_s. &
 \end{array}$$

A manifold which admits such a tower of (principal) bundles is called a **s -step iterated (principal) bundle**. We say that the structure of iterated (principal) bundle is induced by a filtration $\mathcal{S}^k \mathfrak{g}$ in $\mathfrak{g}$ if we have a tower of (principal) bundles as above such that F_i is (the quotient by the lattice of) the Lie-group corresponding to $\mathcal{S}^i \mathfrak{g} / \mathcal{S}^{i-1} \mathfrak{g}$ and G_i is (the quotient by the lattice of the) Lie-group associated to the Lie-algebra $\mathfrak{g} / \mathcal{S}^{i-1} \mathfrak{g}$.

1.2.2. *The complex geometry of the universal covering G .* To study the complex geometry of $(\mathfrak{g}, J)$ we need to study the interplay of $\mathfrak{g}$ and J . The descending central series associated to J is defined by

$$\mathcal{C}_J^i \mathfrak{g} := \mathcal{C}^i \mathfrak{g} + J\mathcal{C}^i \mathfrak{g}$$

and it is known to have the following properties:

- (i) $\mathcal{C}_J^i \mathfrak{g}$ is a J -invariant subalgebra of $\mathfrak{g}$ and an ideal in $\mathcal{C}_J^{i-1} \mathfrak{g}$ [CF01].
- (ii) We have always $\mathcal{C}^i \mathfrak{g} \subset \mathcal{C}_J^i \mathfrak{g}$ but in general inclusion can be strict. Nevertheless we have always $\mathcal{C}_J^1 \mathfrak{g} \neq \mathfrak{g}$. ([Sal01], Theorem. 1.3).
- (iii) There is always a holomorphic fibration of G over the abelian Lie-group (vector space) $\mathfrak{g} / \mathcal{C}_J^1 \mathfrak{g}$ whose typical fibre is the simply connected Lie-group H associated to $\mathcal{C}_J^1 \mathfrak{g}$ with the left-invariant complex structure induced by the restriction of J . This is a real principal H -bundle but in general H will not be a complex Lie-group so there is no way to speak of a holomorphic principal bundle. One can iterate this process to find an iterated holomorphic bundle structure on (G, J) .

The ascending central series associated to J (minimal torus bundle series) is defined by

$$\mathcal{T}^0 \mathfrak{g} := 0, \quad \mathcal{T}^{i+1} \mathfrak{g} := \{x \in \mathfrak{g} \mid [x, \mathfrak{g}] \subset \mathcal{T}^i \mathfrak{g} \text{ and } [Jx, \mathfrak{g}] \subset \mathcal{T}^i \mathfrak{g}\}$$

Every subspace $\mathcal{T}^i \mathfrak{g}$ is J -invariant and an ideal of $\mathfrak{g}$. If there is a k such that $\mathcal{T}^k \mathfrak{g} = \mathfrak{g}$ then the complex structure J is called *nilpotent*. We have $\mathcal{T}^i \mathfrak{g} \subset \mathcal{Z}^i \mathfrak{g}$ for all i and if $\mathcal{T}^i \mathfrak{g} = \mathcal{Z}^i \mathfrak{g}$ then $\mathcal{T}^{i+1} \mathfrak{g}$ is the largest complex subspace of $\mathfrak{g}$ contained in $\mathcal{Z}^{i+1} \mathfrak{g}$ ([CFGU00], Lem. 3).

Not every complex structure is nilpotent, indeed there are examples such that $\mathcal{T}^i \mathfrak{g} = 0$ for all i . In general, if $\mathfrak{g}$ is s -step nilpotent and the complex structure is nilpotent then we have $\mathcal{T}^k \mathfrak{g} = \mathfrak{g}$ for some $k \geq s$ and strict inequality is possible.

Cordero, Fernandez, Gray and Ugarte showed the following [CFGU00]:

Proposition 1.4 — *Let $(\mathfrak{g}, J)$ be a nilpotent Lie-algebra with complex structure. Then J is nilpotent if and only if the associated simply connected Lie-group G has the structure of a k -step iterated principal $\mathbb{C}^{n_i}$ -bundle where k is the smallest integer such that $\mathcal{T}^k \mathfrak{g} := \mathfrak{g}$. The structure of iterated bundle is induced by the filtration $(\mathcal{T}^i \mathfrak{g})$ on the Lie-algebra.*

This bundle structure does not induce a bundle structure on compact quotients of G in general.

1.2.3. *The complex geometry of $M = \Gamma \backslash G$.* We will now address the question of the compatibility of the lattice $\Gamma \subset G$ with the other two structures $\mathfrak{g}$ and J . Most of the cited results originate from the work of Malcev [Mal51].

Definition 1.5 — Let $\mathfrak{g}$ be a nilpotent Lie-algebra. A **rational structure** for $\mathfrak{g}$ is a subalgebra $\mathfrak{g}_{\mathbb{Q}}$ defined over $\mathbb{Q}$ such that $\mathfrak{g}_{\mathbb{Q}} \otimes \mathbb{R} = \mathfrak{g}$.

A subalgebra $\mathfrak{h} \subset \mathfrak{g}$ is said to be rational with respect to a given rational structure $\mathfrak{g}_{\mathbb{Q}}$ if $\mathfrak{h}_{\mathbb{Q}} := \mathfrak{h} \cap \mathfrak{g}_{\mathbb{Q}}$ is a rational structure for $\mathfrak{h}$.

If Γ is a lattice in the corresponding simply connected Lie-group G then its associated rational structure is given by the $\mathbb{Q}$ -span of $\log \Gamma$. A rational subspace with respect to this structure is called **Γ -rational**.

A complex structure J on $\mathfrak{g}$ is called rational if it maps $\mathfrak{g}_{\mathbb{Q}}$ to itself; if $\mathfrak{g}_{\mathbb{Q}}$ is induced by a lattice Γ in the corresponding Lie-group J is called **Γ -rational**.

Remark 1.6 — The lattice and the rational structure are closely related:

- (i) The rational structure associated to Γ is in fact a rational structure ([CG90], p. 204) and there exists a lattice in a nilpotent simply connected Lie-group G if and only if the corresponding Lie-algebra admits a rational structure.
- (ii) After passing to a subgroup of finite index one can always assume that $\log(\Gamma) \subset \mathfrak{g}$ is a lattice, i.e., an additive subgroup which is closed under the Lie bracket. Geometrically this corresponds to a finite covering.
- (iii) If two lattices $\Gamma_1, \Gamma_2 \subset G$ generate the same rational structure then the lattice $\Gamma_1 \cap \Gamma_2$ is of finite index in both of them ([CG90], Theorem 5.1.12, p. 205). Geometrically this corresponds to an étale correspondence

$$\begin{array}{ccc} & (\Gamma_1 \cap \Gamma_2) \backslash G & \\ \swarrow & & \searrow \\ \Gamma_1 \backslash G & & \Gamma_2 \backslash G. \end{array}$$

- (iv) The subalgebras $\mathcal{C}^i \mathfrak{g}$ and $\mathcal{Z}^i \mathfrak{g}$ as defined above are always rational subalgebras ([CG90], p. 208). Thus the filtration $(\mathcal{Z}^i \mathfrak{g})$ yields a structure of real analytic iterated principal torus bundle on $\Gamma \backslash G$.

Lemma 1.7 — If in the situation of Lemma 1.3 $\Gamma \subset G$ is a lattice then π induces a fibration on the compact nilmanifold $M = \Gamma \backslash G$ if and only if $\Gamma \cap H$ is a lattice in H if and only if the associated subalgebra $\mathfrak{h} \subset \mathfrak{g}$ is Γ -rational.

In particular M fibres as a principal holomorphic torus bundle $\pi : M \rightarrow M'$ over some nilmanifold M' if and only if there is a J -invariant, Γ -rational subspace contained in the centre $\mathcal{Z} \mathfrak{g}$ of $\mathfrak{g}$.

Proof. The first part is [CG90], Theorem. 5.1.11, p. 204 and Lem 5.1.4, p. 196. The second assertion then follows. $\square$

To describe effectively the how the geometry of the nilmanifold is encoded in the Lie-algebra we need

Definition 1.8 — Let $\mathfrak{g}$ be a nilpotent Lie-algebra with rational structure $\mathfrak{g}_{\mathbb{Q}}$. We call an ascending filtration $(\mathcal{S}^i \mathfrak{g})_{i=0, \dots, t}$ on $\mathfrak{g}$ a **torus bundle series**

for a complex structure J if for all $i = 1 \dots, t$

- (a) $\mathcal{S}^i \mathfrak{g}$ is rational with respect to $\mathfrak{g}_{\mathbb{Q}}$ and an ideal in $\mathcal{S}^{i+1} \mathfrak{g}$,
- (b) $J\mathcal{S}^i \mathfrak{g} = \mathcal{S}^i \mathfrak{g}$,
- (c) $\mathcal{S}^{i+1} \mathfrak{g} / \mathcal{S}^i \mathfrak{g}$ is an abelian .

If in addition

- (c') $\mathcal{S}^{i+1} \mathfrak{g} / \mathcal{S}^i \mathfrak{g} \subset \mathcal{Z}(\mathfrak{g} / \mathcal{S}^i \mathfrak{g})$,

then $(\mathcal{S}^i \mathfrak{g})_{i=0, \dots, t}$ is called a **principal torus bundle series**.

An ascending filtration $(\mathcal{S}^i \mathfrak{g})_{i=0, \dots, t}$ on $\mathfrak{g}$ is said to be a **stable torus bundle series (STBS)** for $\mathfrak{g}$, if $(\mathcal{S}^i \mathfrak{g})_{i=0, \dots, t}$ is a torus bundle series for every complex structure J and every rational structure $\mathfrak{g}_{\mathbb{Q}}$ in $\mathfrak{g}$. If also condition (c') holds then it is called a **stable principal torus bundle series (SPTBS)**.

Note that we always refer to complex torus bundles and not to real torus bundles. The rationality condition holds for example if every subspace in the series can be described by the subspaces of the ascending and descending central series.

Remark 1.9 — If $M = (\mathfrak{g}, J, \Gamma)$ is a nilmanifold and $\mathfrak{g}$ admits a torus bundle series $(\mathcal{S}^i \mathfrak{g})_{i=0, \dots, t}$ compatible with J and Γ then M can be viewed as an iterated bundle in the following sense: every successive quotient $0 \rightarrow \mathcal{S}^i \mathfrak{g} \rightarrow \mathcal{S}^{i+1} \mathfrak{g} \rightarrow \mathcal{S}^{i+1} \mathfrak{g} / \mathcal{S}^i \mathfrak{g} \rightarrow 0$ induces a holomorphic fibration over a torus with fibre a nilmanifold with left-invariant complex structure. There is not necessarily a iterated torus bundle structure as discussed before since we did not restrict to the case where $\mathcal{S}^{i+1} \mathfrak{g} / \mathcal{S}^i \mathfrak{g}$ is an ideal in $\mathfrak{g} / \mathcal{S}^i \mathfrak{g}$; the map $\mathfrak{g} / \mathcal{S}^i \mathfrak{g} \rightarrow \mathfrak{g} / \mathcal{S}^{i+1} \mathfrak{g}$ might not induce a fibration on the nilmanifold. If J is Γ -rational then the adapted descending central series $\mathcal{C}_j^i \mathfrak{g}$ gives such an example.

The most interesting case is when $(\mathcal{S}^i \mathfrak{g})_{i=0, \dots, t}$ is a SPTBS for $\mathfrak{g}$. Condition (c') implies that $\mathcal{S}^{i+1} \mathfrak{g} / \mathcal{S}^i \mathfrak{g}$ is an ideal in $\mathfrak{g} / \mathcal{S}^i \mathfrak{g}$ so that the filtration $(\mathcal{S}^i \mathfrak{g})_{i=0, \dots, t}$ induces a real analytic iterated principal torus bundle structure on M :

$$\begin{array}{ccc}
 T_1 \hookrightarrow & M_1 = M & \\
 & \downarrow \pi_1 & \\
 T_2 \hookrightarrow & M_2 & \\
 & \downarrow \pi_2 & \\
 & \vdots & \\
 & \downarrow & \\
 T_{s-1} \hookrightarrow & M_{t-1} & \\
 & \downarrow \pi_{t-1} & \\
 & M_t = T_t. &
 \end{array}$$

By definition the maps π_i are holomorphic with respect to any left-invariant complex structure J because the differentials at the identity are given by the complex linear maps

$$\mathcal{S}^i \mathfrak{g} / \mathcal{S}^{i-1} \rightarrow \mathfrak{g} / \mathcal{S}^{i-1}.$$

A SPTBS gives in some sense the appropriate description of the complex geometry of the corresponding nilmanifolds.

In general, a STBS, let alone a SPTBS, will not exist for a nilpotent Lie-algebra but it is not as uncommon as one might think at first sight. In real dimension 6 the Lie-algebra $\mathfrak{h}_7$ described in Example 1.14 is the only one not to admit at least a STBS.

1.2.4. Abelian and complex parallelisable structures. There are two extremal types of complex structures that can occur in a nilpotent Lie-algebra with complex structure $(\mathfrak{g}, J)$. If $[\mathfrak{g}^{1,0}, \mathfrak{g}^{0,1}] = 0$ then we have $[Jx, y] = J[x, y]$ for all $x, y \in \mathfrak{g}$; thus the Lie-bracket is $\mathbb{C}$ -linear and $(\mathfrak{g}, J)$ is a complex Lie-algebra. Nilmanifolds of type $(\mathfrak{g}, J)$ are then complex parallelisable. These have been studied in detail by Winkelmann [Win98] and have very special (arithmetic) properties. Both the descending and the ascending central series are principal torus bundle series (which do not need to be stable). Their small deformations have been described in [Rol08a].

If $(\mathfrak{g}, J)$ satisfies the complementary condition $[\mathfrak{g}^{1,0}, \mathfrak{g}^{1,0}] = 0$ then the complex structure is called abelian and for such nilmanifolds the ascending central series is a principal torus bundle series. These structure were extensively studied by Salamon, McLaughlin, Pedersen, Poon, Console, Fino e.a. for example in [CFP06] and [MPPS06].

1.3. Dolbeault cohomology of nilmanifolds and small deformations.

Let $M_J = (\mathfrak{g}, J, \Gamma)$ be a nilmanifold with left-invariant complex structure. In this section we want to describe how cohomology and deformations of M_J are governed by $(\mathfrak{g}, J)$.

By a theorem of Nomizu [Nom54] the de Rham cohomology of a nilmanifold can be calculated using left-invariant differential forms and is isomorphic to the cohomology of the complex

$$0 \rightarrow \mathfrak{g}^* \xrightarrow{d} \Lambda^2 \mathfrak{g}^* \xrightarrow{d} \Lambda^3 \mathfrak{g}^* \xrightarrow{d} \dots,$$

which also calculates the Lie-algebra cohomology of $\mathfrak{g}$.

The question if the Dolbeault cohomology of nilmanifolds with left-invariant complex structure can be calculated using invariant differential forms has been addressed by Console and Fino in [CF01] and Cordero, Fernandez, Gray and Ugarte in [CFGU00]: consider in the Dolbeault-complex for $\Omega_{M_J}^p$ the subcomplex of left-invariant forms

$$0 \rightarrow \Lambda^p \mathfrak{g}^{*1,0} \xrightarrow{\bar{\partial}} \Lambda^p \mathfrak{g}^{*1,0} \otimes \mathfrak{g}^{*0,1} \xrightarrow{\bar{\partial}} \Lambda^p \mathfrak{g}^{*1,0} \otimes \Lambda^2 \mathfrak{g}^{*0,1} \xrightarrow{\bar{\partial}} \dots$$

and denote by $H^{p,q}(\mathfrak{g}, J)$ its q -th cohomology group. Using our definitions their results read as follows:

Theorem 1.10 — *Let $M_J = (\mathfrak{g}, J, \Gamma)$ be a nilmanifold with left-invariant complex structure. Then the inclusion*

$$(2) \quad \iota_J : H^{p,q}(\mathfrak{g}, J) \rightarrow H^{p,q}(M_J)$$

is an isomorphism if

- (i) The complex structure J is bi-invariant, G is a complex Lie-group and M_J is complex parallelisable [Sak76].
- (ii) The complex manifold M_J is an iterated principal holomorphic torus bundle [CFGU00].
- (iii) The complex structure J is Γ -rational. ([CF01], Theorem B)
- (iv) The complex structure J is abelian [CF01].
- (v) $\mathfrak{g}$ admits a torus bundle series for J compatible with the rational structure induced by Γ .

Moreover there is a dense open subset U of the space $\mathcal{C}(\mathfrak{g})$ of all left-invariant complex structures on M such that ι is an isomorphism for all $J \in U$ ([CF01], Theorem A).

Statement (v) follows immediately from the the proof of Theorem B in [CF01]. The idea of their proof is the following: as long as M_J can be given a structure of iterated bundle with a good control over the cohomology of the base and of the fibre, one can use the Borel spectral sequence for Dolbeault cohomology in order to get an inductive proof. If the complex structure is Γ -rational then the descending central series adapted to J induces such an iterated fibration which yields the result on a dense subset of the space of invariant complex structures; Console and Fino then show that the property “The map ι_J is an isomorphism.” is stable under small deformations.

It is an open question if ι_J is an isomorphism for every nilmanifold with left-invariant complex structure.

Generalising results on deformations of abelian complex structures [MPPS06, CFP06] we proved the following

Theorem 1.11 ([Rol08b], Theorem 2.6) — *Let $M_J = (\mathfrak{g}, J, \Gamma)$ be a nilmanifold with left-invariant complex structure such that*

$$\iota : H^{1,q}(\mathfrak{g}, J, \mathbb{C}) \rightarrow H^{1,q}(M_J)$$

is an isomorphism for all q . Then all small deformations of the complex structure J are again left-invariant complex structures. More precisely, the Kuranishi family contains only left-invariant complex structures.

Together with Theorem 1.10 (v) this yields

Corollary 1.12 — *Let M_J be a nilmanifold of type $(\mathfrak{g}, \Gamma)$. If $\mathfrak{g}$ admits a stable (principal) torus bundle series then every small deformation of M_J is a nilmanifold with left-invariant complex structure of type $(\mathfrak{g}, \Gamma)$.*

In other words, if M is a real nilmanifold with Lie-algebra $\mathfrak{g}$ and $\mathfrak{g}$ admits a STBS then the subset of left-invariant complex structures is open in the space of all complex structures. This is a first step on our way to prove that in some cases left-invariant complex structures form a connected component of the space of complex structures.

1.4. Examples and Counterexamples. The simplest example of a nilmanifolds with left-invariant complex structure is a complex torus which corresponds to an abelian Lie-algebra. Other well known examples are Kodaira surfaces, which were historically the first manifolds known to admit

both a symplectic structure and a complex structure but no Kähler structure. These are principal bundles of elliptic curves over elliptic curves.

The following list gives an overview over some phenomena that can occur:

- Small deformations of iterated principal bundles do not need to have any torus bundle structure: Example 1.14.
- First examples with SPTBS: Complex tori, Kodaira surfaces, principal bundles over elliptic curve (Example 1.13).
- The minimal principal torus bundle series must not be stable: Example 1.15.
- A Lie-algebra which admits only non-nilpotent complex structures: $\mathfrak{h}_{26}^+$ discussed in Section 4.2.
- A Lie-algebra which admits both nilpotent and non-nilpotent complex structures: Example 3.13, see also [CFP06], Section 6.
- The dimension of the Albanese variety is not constant under small deformations: Example 2.4.

In order to describe the examples we have to explain some notation. Consider a Lie-algebra $\mathfrak{g}$ spanned by a basis $e_1, \dots, e_n$. Then the Lie-bracket is uniquely determined by structure constants a_{ij}^k such that

$$[e_i, e_j] = \sum_{k=1}^n a_{ij}^k e_k$$

satisfying $a_{ij}^k = -a_{ji}^k$ and the relations encoding the Jacobi identity. Let $\langle e^1, \dots, e^n \rangle$ be the dual basis, i.e. $e^i(e_j) = \delta_{ij}$. Then for any $\alpha \in \mathfrak{g}^*$ and $x, y \in \mathfrak{g}$ we define

$$d : \mathfrak{g}^* \rightarrow \Lambda^2 \mathfrak{g}^*, \quad d\alpha(x, y) := -\alpha([x, y])$$

and get a dual description of the Lie-bracket by

$$de^k = - \sum_{i < j} a_{ij}^k e^{ij}$$

where we abbreviate $e^i \wedge e^j = e^{ij}$. The map d induces a map on the exterior algebra $\Lambda^* \mathfrak{g}^*$ and $d^2 = 0$ is equivalent to the Jacobi identity. Giving a 2-step Lie-algebra is the same as giving a surjective alternating bilinear form on $\mathfrak{g}/\mathcal{Z}\mathfrak{g}$ with values in $\mathcal{Z}\mathfrak{g}$; the Jacobi Identity is trivial in this case.

If we write $\mathfrak{h}_2 = (0, 0, 0, 0, 12, 34)$ we mean the following: Let $e_1, \dots, e_6$ be a basis for the Lie-algebra and $e^1, \dots, e^6$ be the dual basis. Then the defining relations for $\mathfrak{h}_2$ are given by

$$de^1 = de^2 = de^3 = de^4 = 0, \quad de^5 = e^{12} := e^1 \wedge e^2, \quad de^6 = e^{34}.$$

Dually, the bracket relations are generated by $[e_1, e_2] = -e_5$ and $[e_3, e_4] = -e_6$. This Lie-algebra will come up again in Section 4.2 when we discuss complex structures on real 6-dimensional Lie-algebras.

We will usually describe a complex structure on a Lie-algebra by specifying an endomorphism J on the basis of $\mathfrak{g}$ such that $J^2 = -id_{\mathfrak{g}}$; if we set $Je_1 = e_2$ then $Je_2 = J^2 e_1 = -e_1$ is tacitly understood. An equivalent way is to give the decomposition $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g}^{1,0} \oplus \mathfrak{g}^{0,1}$ in the $(\pm i)$ -eigenspaces of J .

There is a very simple class of examples to which our results will apply:

Example 1.13 — Let M_J be a non-trivial principal holomorphic torus bundle over an elliptic curve. Regarding M_J as a nilmanifold with left-invariant complex structure with Lie algebra $\mathfrak{g}$, the fibration corresponds to a central extension

$$0 \rightarrow \mathcal{T}^1 \mathfrak{g} \rightarrow \mathfrak{g} \rightarrow \mathfrak{e} \rightarrow 0$$

where $\mathcal{T}^1 \mathfrak{g}$ is the $(2n-2)$ -dimensional real, J -invariant subspace corresponding to the fibres of the bundle. In particular the centre of $\mathfrak{g}$ has dimension at least $2n-2$ whence, since M is not a torus, we have $\mathcal{Z}\mathfrak{g} = \mathcal{T}^1 \mathfrak{g}$. There is only one Lie-algebra with 2-codimensional centre so $\mathfrak{g} \cong (0, 0, 0, \dots, 0, 12)$. We will see in Proposition 3.6 that

$$\mathfrak{g} \supset \mathcal{Z}\mathfrak{g} \supset 0$$

is a SPTBS for $\mathfrak{g}$.

Example 1.14 — A nice behaviour of the complex structure on the universal covering G may not be sufficient to get a good description the manifold $M = \Gamma \backslash G$.

Consider the Lie-group H_7 whose Lie-algebra is $\mathfrak{h}_7 = (0, 0, 0, 12, 13, 23)$ with basis $e_1, \dots, e_6$. The vectors $e_4, \dots, e_6$ span the centre $\mathcal{Z}^1 \mathfrak{h}_7 = \mathcal{C}^1 \mathfrak{h}_7$.

Let $\Gamma \subset H_7$ be the lattice generated the elements $\exp(e_k)$ and consider the nilmanifold $M = \Gamma \backslash H_7$, which – as a real manifold – can be regarded as a real principal torus bundle with fibre and base a 3-dimensional torus.

For every real number λ we give a left-invariant complex structure J_λ on M by specifying a basis for the space of $(1, 0)$ -vectors:

$$(\mathfrak{h}_7^{1,0})_\lambda := \langle X_1 = e_1 - ie_2, X_2^\lambda = e_3 - i(e_4 - \lambda e_1), X_3^\lambda = -e_5 - \lambda e_4 + ie_6 \rangle$$

One can check that $[X_1, X_2^\lambda] = X_3^\lambda$ and, since X_3^λ is contained in the centre, the complex structure is integrable. The largest complex subspace of the centre is spanned by the real and imaginary part of X_3^λ since the centre has real dimension three.

The simply connected Lie-group H_7 has a filtration by subgroups induced by the filtration

$$\mathfrak{h}_7 \supset V_1 = \langle \lambda e_2 + e_3, e_4, \operatorname{Im}(X_3^\lambda), \operatorname{Re}(X_3^\lambda) \rangle \supset V_2 = \langle \operatorname{Im}(X_3^\lambda), \operatorname{Re}(X_3^\lambda) \rangle \supset 0$$

on the Lie-algebra and, since all these are J invariant, H_7 has the structure of a tower of principal holomorphic bundles with fibre $\mathbb{C}$, i.e., the complex structure is nilpotent. In fact, using the results of [Uga04] a simple calculation shows that every complex structure on $\mathfrak{h}_7$ is equivalent to J_0 .

Now we take the compatibility with the lattice into account. The rational structure induced by Γ coincides with the $\mathbb{Q}$ -algebra generated by the basis vectors e_k . Therefore $V_j \cap \Gamma$ is a lattice in V_j if and only if λ is rational. That is, for $\lambda \notin \mathbb{Q}$ the structure of iterated holomorphic principal bundle on H_7 does not descend to the quotient $\Gamma \backslash H_7$.

Example 1.15 — The aim of this example is to show that the minimal torus bundle series $\mathcal{T}^i \mathfrak{g}$ might not yield the appropriate geometric description of an associated nilmanifold.

We consider the Lie-algebra given by $\mathfrak{g} = (0, 0, 0, 0, 0, 12, 13)$ with the lattice $\Gamma \subset G$ generated by the images of the basis vectors e_k . Let $\langle z_1, z_2, z_3 \rangle$

be any basis of the subspace spanned by $e_4, \dots, e_6$. We define a complex structure by declaring

$$\begin{aligned} J e_2 &= e_3, & J e_7 &= e_8, \\ J z_1 &= e_1, & J z_2 &= z_3. \end{aligned}$$

A short calculation shows that J is integrable and that $\mathcal{T}^1 \mathfrak{g} = \langle z_2, z_3, e_7, e_8 \rangle$ is the largest complex subspace of the centre. But the z_i were quite arbitrarily chosen and hence $\mathcal{T}^1 \mathfrak{g}$ is not a Γ -rational subspace in general, which means that it does not tell us a lot about the geometry of the compact manifolds $M = \Gamma \backslash G$. But on the other hand $\mathcal{C}^1 \mathfrak{g} = \langle e_7, e_8 \rangle$ is a Γ -rational, J -invariant subspace. This is no coincidence: we will prove in Proposition 3.8 that

$$0 \subset \mathcal{C}^1 \mathfrak{g} \subset \mathfrak{g}$$

is a SPTBS for $\mathfrak{g}$ which means that we can describe M as a principal bundle of elliptic curves over a complex 3-torus for any complex structure on $\mathfrak{g}$.

2. ALBANESE-QUOTIENTS AND DEFORMATIONS IN THE LARGE

In this section Δ will denote a small 1-dimensional disc centred in 0. In [Cat04] Catanese proved that any deformation in the large of a complex torus is again a complex torus by analysing the Albanese map: he studies families $\mathcal{X} \rightarrow \Delta$ with general fibre a complex torus. In a first step he shows that also the special fibre $\mathcal{X}_0$ has a surjective Albanese map to a complex torus of the same dimension. By proving that this map has to be biholomorphic he concludes that the special fibre is a complex torus. We will try to generalise his method to the case where the Albanese map is not so well behaved.

2.1. Definitions and results. We need to recall some definitions. Let X be a compact, complex manifold. By Kodaira's Lemma ([Cat04], Lemma 2.2) we have an inclusion

$$H^0(X, d\mathcal{O}_X) \oplus \overline{H^0(X, d\mathcal{O}_X)} \subset H_{dR}^1(X, \mathbb{C}).$$

Definition 2.1 — *The **Albanese variety** of X is the abelian, complex Lie-group $\text{Alb}(X)$ defined as the quotient of $H^0(X, d\mathcal{O}_X)^*$ by the minimal, closed, complex subgroup containing the image of $H_1(X, \mathbb{Z})$ under the map*

$$H_1(X, \mathbb{Z}) \rightarrow H_1(X, \mathbb{C}) \rightarrow H^0(X, d\mathcal{O}_X)^*.$$

*The Albanese variety is called **very good** if the image of $H_1(X, \mathbb{Z})$ is a lattice in $H^0(X, d\mathcal{O}_X)^*$.*

The Albanese map $\alpha_X : X \rightarrow \text{Alb}(X)$ is given by integration of closed forms along paths starting from a fixed base point.

Remark 2.2 — Jörg Winkelmann brought to our attention that the Albanese variety is always compact and hence $\text{Alb}(X)$ is very good if and only if the image of $H_1(X, \mathbb{Z})$ is discrete in $H^0(X, d\mathcal{O}_X)^*$.

If X satisfies the weak 1-Hodge property which means

$$H_{dR}^1(X, \mathbb{C}) = H^0(X, d\mathcal{O}_X) \oplus \overline{H^0(X, d\mathcal{O}_X)} = H^0(X, d\mathcal{O}_X) \oplus H^1(X, \mathcal{O}_X)$$

then X has a very good Albanese variety; in particular this holds if X is kählerian.

Catanese studied the behaviour of the Albanese map with respect to deformation and obtained for example the following

Proposition 2.3 ([Cat04], Corollary 2.5) — *Assume that $\{X_t\}_{t \in \Delta}$ is a 1-parameter family of compact complex manifolds over the unit disc such that there is a sequence $t_\nu \rightarrow 0$ with X_{t_ν} satisfying the weak 1-Hodge property, and moreover such that the image of X_{t_ν} under the Albanese map has dimension a . Then also the central fibre has a very good Albanese map and Albanese dimension a .*

In the following we want to explain how to generalise this result if the Albanese variety is not necessarily very good; the right way to do so is best explained via the following example.

Example 2.4 — Let $X = (\mathfrak{g}, J, \Gamma)$ be a principal holomorphic torus bundle over an elliptic curve as in Example 1.13. Then there is a basis $e_1, e_2, z_1, \dots, z_{2k}$ of $\mathfrak{g}$ such that the only non-trivial Lie-bracket relation is $[e_1, e_2] = z_{2k}$ and $\mathcal{C}^1 \mathfrak{g}$ is spanned by z_{2k} . The centre $\mathcal{Z} \mathfrak{g} = \langle z_1, \dots, z_{2k} \rangle$ corresponds to the fibres of the bundle and is always J -invariant. With respect to the dual basis $e^1, \dots, z^{2k}$ the differential $d : \mathfrak{g}^* \rightarrow \Lambda^2 \mathfrak{g}^*$ is given by

$$dz^{2k} = -e^1 \wedge e^2, de^1 = de^2 = dz^1 = \dots = dz^{2k-1} = 0$$

and, by the description of cohomology given in Section 1.3, we see that

$$H^1(X, \mathbb{C}) = \{\alpha \in \mathfrak{g}_{\mathbb{C}}^* \mid d\alpha = 0\}$$

has dimension $2k + 1$. Thus X can never satisfy the weak 1-Hodge property.

In order to determine the Albanese Variety of X we have to take the rational structure on $\mathfrak{g}$ induced by the lattice Γ into account; on the quotient $H_1(X, \mathbb{R}) = \mathfrak{g} / \mathcal{C}^1 \mathfrak{g}$ it maps to the rational structure induced by $H_1(X, \mathbb{Z})$. Let W be the smallest J -invariant, Γ -rational subspace of $\mathfrak{g}$ which contains $\mathcal{C}^1 \mathfrak{g}$. Since the centre is J -invariant and Γ -rational we have

$$2 \leq \dim_{\mathbb{R}} W \leq \dim_{\mathbb{R}} \mathcal{Z} \mathfrak{g} = 2k.$$

The space of closed holomorphic 1-forms is

$$\begin{aligned} H^0(X, d\mathcal{O}_X) &= \{\alpha \in \mathfrak{g}^{*1,0} \mid d\alpha = 0\} \\ &= \{\alpha \in \mathfrak{g}^{*1,0} \mid \bar{\partial}\alpha = 0\} \\ &= \text{Ann}((\mathcal{C}_J^1 \mathfrak{g}) \otimes \mathbb{C}) \cap \mathfrak{g}^{*1,0}, \end{aligned}$$

hence the kernel of the surjection $p : H_1(X, \mathbb{R}) \twoheadrightarrow H^0(X, d\mathcal{O}_X)^*$ is $\mathcal{C}_J^1 \mathfrak{g} / \mathcal{C}^1 \mathfrak{g}$ yielding

$$\text{Alb}(X) = \frac{H^0(X, d\mathcal{O}_X)^* / p(W)}{\text{im}(H_1(X, \mathbb{Z}) \rightarrow H^0(X, d\mathcal{O}_X)^* / p(W))}.$$

The dimension of the Albanese variety can vary with the complex structure in the range

$$1 \leq \dim \text{Alb}(X) = \frac{\dim_{\mathbb{R}} \mathfrak{g} - \dim_{\mathbb{R}} W}{2} \leq k = h^0(X, d\mathcal{O}_X).$$

But in some sense all the ugly phenomena happen inside $\mathcal{Z} \mathfrak{g}$. Indeed, if we replace W by $\mathcal{Z} \mathfrak{g}$ in the above construction we get a map to an elliptic curve such that the corresponding map in cohomology is independent of the complex structure; we have found a way to reconstruct our map π :

$X \rightarrow E$ from cohomological data which does not depend on the particular left-invariant complex structure.

Thus, even if the Albanese map is not well behaved, we might find some subspace in $H_1(X, \mathbb{R})$, not depending on the complex structure, which yields a holomorphic map to a compact torus. We will now formalise this idea.

Let $V \subsetneq H_1(X, \mathbb{Q})$ be a proper subspace and let

$$U := \text{Ann}(V_{\mathbb{C}}) \cap H^0(X, d\mathcal{O}_X)$$

where the annihilator is taken with respect to the pairing between $H^1(X, \mathbb{C})$ and $H_1(X, \mathbb{C})$. If $\text{Ann}(V_{\mathbb{C}}) = U \oplus \bar{U}$ then the map $H_1(X, \mathbb{R})/V_{\mathbb{R}} \xrightarrow{\cong} U^*$ is an isomorphism of real vector spaces.

Since V is a rational subspace, $H_1(X, \mathbb{Z})/(H_1(X, \mathbb{Z}) \cap V)$ maps to a full lattice $\Lambda \subset U^*$. This yields a complex torus U^*/Λ which is a quotient of the Albanese variety $\text{Alb}(X)$, the projection being induced by the inclusion $U \hookrightarrow H^0(X, d\mathcal{O}_X)$.

We have shown that the following is well defined and yields in fact a compact, complex torus:

Definition 2.5 — Let $V \subsetneq H_1(X, \mathbb{Q})$ be a proper subspace such that

$$\text{Ann}(V_{\mathbb{C}}) = U \oplus \bar{U}$$

where $U := \text{Ann}(V_{\mathbb{C}}) \cap H^0(X, d\mathcal{O}_X)$. Then we call

$$Q\text{Alb}_V(X) := U^*/\Lambda$$

the **very good Albanese-Quotient** of X associated to V , where Λ is the image of the lattice $H_1(X, \mathbb{Z})/(H_1(X, \mathbb{Z}) \cap V)$ under the map $H_1(X, \mathbb{R})/V_{\mathbb{R}} \rightarrow U^*$.

Remark 2.6 — The above condition on V is satisfied if and only if the kernel of the composition map

$$\phi : H^0(X, d\mathcal{O}_X) \rightarrow H^1(X, \mathbb{C}) \rightarrow H^1(X, \mathbb{C})/\text{Ann}(V_{\mathbb{C}})$$

has complex dimension $q = \frac{\dim_{\mathbb{Q}}(V)}{2}$. This is in fact the maximal possible dimension since $V_{\mathbb{C}}$ is defined over $\mathbb{R}$ while on the other hand

$$H^0(X, d\mathcal{O}_X) \cap \overline{H^0(X, d\mathcal{O}_X)} = 0.$$

Another way to look at this is the following: let

$$W := (H^0(X, d\mathcal{O}_X) \oplus \overline{H^0(X, d\mathcal{O}_X)}) \cap H^1(X, \mathbb{R}).$$

The decomposition $W_{\mathbb{C}} = H^0(X, d\mathcal{O}_X) \oplus \overline{H^0(X, d\mathcal{O}_X)}$ defines a complex structure on W and the above conditions on V are equivalent to $\text{Ann}(V_{\mathbb{C}})$ being a complex subspace of W with respect to this complex structure.

Remark 2.7 — Integration over the closed forms which are contained in $\text{Ann}(V_{\mathbb{C}}) \cap H^0(X, d\mathcal{O}_X)$ gives us a holomorphic map $q\alpha_V$ which factors through the Albanese map:

$$\begin{array}{ccc} X & \xrightarrow{\alpha} & \text{Alb}(X) \\ & \searrow q\alpha_V & \downarrow \\ & & Q\text{Alb}_V(X). \end{array}$$

Note that $Q\text{Alb}_V(X)$ has positive dimension by definition since we assumed $V \neq H_1(X, \mathbb{Q})$.

Example 2.8 — Let $M = (\mathfrak{g}, J, \Gamma)$ be a 2-step nilmanifold such that the centre $\mathcal{Z}\mathfrak{g}$ is J -invariant. By Nomizu's theorem we have $H_1(M, \mathbb{R}) \cong \mathfrak{g}/[\mathfrak{g}, \mathfrak{g}]$ and $V_{\mathbb{R}} := \mathcal{Z}\mathfrak{g}/[\mathfrak{g}, \mathfrak{g}]$ is a Γ -rational subspace. Then $V_{\mathbb{Q}}$ induces a very good Albanese-Quotient and $q\alpha_V$ coincides with the principal holomorphic bundle map induced by the filtration $\mathfrak{g} \supset \mathcal{Z}\mathfrak{g} \supset 0$. In particular this applies to the principal holomorphic torus bundle over an elliptic curve discussed in Example 2.4 above.

Our goal is to study such locally trivial, smooth, holomorphic fibrations. Hence we define

Definition 2.9 — Let $\mathcal{C}$ be a class of compact, complex manifolds and let X be any compact, complex manifold. We say that a subspace $V \subsetneq H_1(X, \mathbb{Q})$ is **$\mathcal{C}$ -fibering** on X if V induces a very good Albanese-Quotient and the map

$$q\alpha : X \rightarrow Q\text{Alb}_V(X)$$

is a locally trivial, smooth, holomorphic fibration with fibres in $\mathcal{C}$.

A class $\mathcal{C}$ of compact complex manifolds is called a **good fibre class** if the following conditions hold

- (i) The class $\mathcal{C}$ is closed under holomorphic 1-parameter limits, i.e., if we have a smooth family over the unit disc and a sequence $(t_\nu)_{\nu \in \mathbb{N}}$ converging to 0 such that the fibres over t_ν are in $\mathcal{C}$ then also the central fibre is in $\mathcal{C}$.
- (ii) One of the following conditions holds:
 - (a) There is a coarse moduli space $\mathfrak{M}_{\mathcal{C}}$ for manifolds in class $\mathcal{C}$ which is Hausdorff.
 - (b) For every manifold $Y \in \mathcal{C}$ there is a local moduli space, i.e., the Kuranishi family of Y is universal.
 - (c) $h^1(F, \Theta_F)$ is constant on the connected components of $\mathcal{C}$.

So in the above example the subspace V is torus-fibering.

Remark 2.10 — The condition (a) is very strong but it is worth including since it applies to Kodaira surfaces. Indeed these and complex tori are our first examples of good fibre classes.

That every deformation in the large of a complex torus is again a complex torus has been proved by Catanese in [Cat02] and we have $h^1(T, \Theta_T) = \dim(T)^2$ for any Torus T (for example by Section 1.3).

Borcea proved in [Bor84] that the moduli space of Kodaira surfaces may be identified with the product of the complex plane by the punctured disc which is Hausdorff. That every deformation in the large is again a Kodaira surface has been proved by the author in [Rol07b].

The main result of this section is

Theorem 2.11 — Let $\mathcal{C}$ be a good fibre class and $\pi : \mathcal{X} \rightarrow \Delta$ be a smooth family of compact, complex manifolds. Let $(t_\nu)_{\nu \in \mathbb{N}}$ be a sequence in Δ converging to zero.

If there is a subspace $V \subsetneq H_1(\mathcal{X}_0, \mathbb{Q})$ which is $\mathcal{C}$ -fibering on $\mathcal{X}_{t_\nu}$ for all ν then V is $\mathcal{C}$ -fibering on $\mathcal{X}_0$.

The proof will be given in the next subsection. We will first show how this can be applied to nilmanifolds

Theorem 2.12 — *Let G be a simply connected nilpotent Lie-group with Lie-algebra $\mathfrak{g}$ and let $\Gamma \subset G$ be a lattice such that the following holds:*

- (i) $\mathfrak{g}$ admits a STBS $(\mathcal{S}^i \mathfrak{g})_{i=0,\dots,t}$ (cf. Definition 1.8).
- (ii) The nilmanifolds with left-invariant complex structure of the type $(\mathcal{S}^{t-1} \mathfrak{g}, \Gamma \cap \exp(\mathcal{S}^{t-1} \mathfrak{g}))$ constitute a good fibre class.

Then any deformation in the large of a nilmanifold with left-invariant complex structure $M_J = (\mathfrak{g}, J, \Gamma)$ is again of type $(\mathfrak{g}, \Gamma)$.

Proof. We will use the following:

Lemma 2.13 ([Cat04], Lemma 2.8) — *Let $\mathcal{B}$ be a connected complex analytic space and $\mathcal{B}'$ a non-empty, open subset of $\mathcal{B}$ such that $\mathcal{B}'$ is closed for holomorphic 1-parameter limits (i.e., given any holomorphic map $f : \Delta \rightarrow \mathcal{B}$, if there is a sequence $(t_\nu)_{\nu \in \mathbb{N}}$ converging to 0 with $f(t_\nu) \in \mathcal{B}'$ then $f(0) \in \mathcal{B}'$). Then $\mathcal{B} = \mathcal{B}'$.*

Now consider any smooth family $\mathcal{X} \rightarrow \mathcal{B}$ over a connected base $\mathcal{B}$ such that for some point $b \in \mathcal{B}$ the fibre $\mathcal{X}_b$ is isomorphic to a nilmanifold $M = (\mathfrak{g}, J, \Gamma)$. Then consider the (non-empty) set

$$\mathcal{B}' := \{t \in \mathcal{B} \mid \mathcal{X}_t \cong M' = (\mathfrak{g}, J', \Gamma) \text{ for some complex structure } J'\}.$$

This set is open in view of our result on small deformations in Corollary 1.12. We have to show that it is closed under holomorphic 1-parameter limits.

Our definition of STBS guarantees that $\Gamma' := \Gamma \cap \exp(\mathcal{S}^{t-1} \mathfrak{g})$ is a lattice. Taking

$$V_{\mathbb{R}} = \mathcal{S}^{t-1} \mathfrak{g} / \mathcal{C}^1 \mathfrak{g} \subset \mathfrak{g} / \mathcal{C}^1 \mathfrak{g} \cong H^1(M, \mathbb{R})$$

we can apply Theorem 2.11 which yields that any 1-parameter limit $\hat{M}$ of manifolds in $\mathcal{B}'$ is a smooth, locally trivial, holomorphic fibration over a torus with fibre a nilmanifold $F = (\mathcal{S}^{t-1} \mathfrak{g}, J, \Gamma')$. The topological structure of the fibre bundle is determined by the fundamental group, which does not change under deformation, and the fundamental group Γ determines G and $\mathfrak{g}$ up to isomorphism ([VGS00], p. 45, corollary 2.8). So also $\hat{M}$ is a nilmanifolds with left-invariant complex structure of the same type.

We have proved that $\hat{M} \in \mathcal{B}'$ hence $\mathcal{B}'$ is closed under holomorphic 1-parameter limits and, by the lemma, $\mathcal{B} = \mathcal{B}'$ which concludes the proof. $\square$

Using Remark 2.10 we see as a first application that every deformation in the large of a principal holomorphic torus bundle over an elliptic curve is again such a principal holomorphic bundle. Further applications will be given in Section 4.

2.2. Proof of Theorem 2.11. We will split the proof of the theorem into several steps.

Catanese showed in [Cat91] that the Albanese dimension is in fact a topological property if X is a Kähler manifold and we review his arguments in our context:

Assume that we have X and $V \subset H_1(X, \mathbb{Q})$ as in Definition 2.5.

Lemma 2.14 — *The dimension of the image of X under the map $q\alpha$ is*

$$d = \max\{m \mid \text{im}(\Lambda^m(\text{Ann}(V_{\mathbb{C}}) \cap H_0(d\mathcal{O}_X)) \rightarrow H^m(X, \mathbb{C})) \neq 0\}.$$

Moreover $q\alpha$ is surjective if and only if $\Lambda^{2k}\text{Ann}(V_{\mathbb{C}}) \neq 0$ in $H^{2k}(X, \mathbb{C})$ where $k = \dim \text{Ann}(V_{\mathbb{C}})$. In particular, the surjectivity of $q\alpha$ is a property which depends only on the topology of X and the subspace V .

Proof. By the definition of $Q\text{Alb}_V(X)$ we have

$$U := q\alpha^*H^0(Q\text{Alb}_V(X), d\mathcal{O}_{Q\text{Alb}_V(X)}) = \text{Ann}(V) \cap H_0(d\mathcal{O}_X)$$

and $q\alpha$ is given by integration over the holomorphic 1-forms in U .

The dimension of its image is, by Sard's theorem, equal to the maximal rank of the differential of the quotient Albanese map and hence equal to d .

It remains to show that $\Lambda^{2k}\text{Ann}(V) \neq 0$ if and only if $\Lambda^k U \neq 0$. But this is clear since our assumptions guarantee that $\text{Ann}(V)$ is contained in $H^0(X, d\mathcal{O}_X) \oplus \overline{H^0(X, d\mathcal{O}_X)}$, hence $\text{Ann}(V)_{\mathbb{C}} = U \oplus \bar{U}$ and $\Lambda^{2k}\text{Ann}(V) = \Lambda^k U \otimes \Lambda^k \bar{U}$. $\square$

In order to analyse how our notion of Albanese-Quotient behaves under deformation we need to introduce some notation.

Let $\pi : \mathcal{X} \rightarrow \Delta$ be a smooth family of compact complex manifolds over the unit disk. We identify $H_1(\mathcal{X}_t, \mathbb{Q})$ with $H_1(\mathcal{X}_0, \mathbb{Q})$.

The sheaves of relative differential forms $\Omega_{\mathcal{X}/\Delta}^k$ are defined by the exact sequence

$$0 \rightarrow \pi^*\Omega_{\Delta}^k \rightarrow \Omega_{\mathcal{X}}^k \rightarrow \Omega_{\mathcal{X}/\Delta}^k \rightarrow 0$$

and we have a $\pi^*\mathcal{O}_{\Delta}$ -linear map

$$d_v : \Omega_{\mathcal{X}/\Delta}^{k-1} \xrightarrow{d} \Omega_{\mathcal{X}}^k \rightarrow \Omega_{\mathcal{X}/\Delta}^k$$

given by differentiation along the fibres. This gives us a complex of sheaves

$$\mathcal{E}^{\bullet} = 0 \rightarrow \mathcal{O}_{\mathcal{X}} \rightarrow \Omega_{\mathcal{X}/\Delta}^1 \rightarrow \Omega_{\mathcal{X}/\Delta}^2 \rightarrow \dots$$

which restricts to the holomorphic de Rham complex on every fibre $\mathcal{X}_t$ of π .

The following useful fact has been extracted from the proof of Lemma 2.4 in [Cat04]. Denote by $\mathbb{C}_t$ the residue field at the point $t \in \Delta$.

Lemma 2.15 — *Let $\pi : \mathcal{X} \rightarrow \Delta$ be a smooth family of compact complex manifolds over a small disk and assume that there is a sequence $(t_{\nu})_{\nu \in \mathbb{N}}$ converging to 0 such that $H^0(\mathcal{X}_{t_{\nu}}, d\mathcal{O}_{\mathcal{X}_{t_{\nu}}})$ has dimension q for all ν .*

Then, after possibly shrinking Δ , there is a rank q locally free subsheaf $\mathcal{H}$ of $H^1(\mathcal{X}_0, \mathbb{C}) \otimes \mathcal{O}_{\Delta}$, the trivial vector bundle on Δ with fibre $H^1(\mathcal{X}_0, \mathbb{C})$, such that

$$\mathcal{H} \otimes \mathbb{C}_t \subset H^0(\mathcal{X}_t, d\mathcal{O}_{\mathcal{X}_t}) \text{ for all } t \in \Delta$$

and equality holds for $t \neq 0$.

Proof. Let $\Delta^* = \Delta \setminus \{0\}$ be the pointed disk. We define a (not necessarily locally free) sheaf $d\mathcal{O}_{\mathcal{X}/\Delta}$ on $\mathcal{X}$ by the exact sequence

$$0 \rightarrow d\mathcal{O}_{\mathcal{X}/\Delta} \rightarrow \Omega_{\mathcal{X}/\Delta}^1 \xrightarrow{d_v} \Omega_{\mathcal{X}/\Delta}^2$$

and push down this sequence to Δ via π_* .

Since the sheaves $\Omega_{\mathcal{X}/\Delta}^k$ are torsion free the same holds for their direct images $\pi_*\Omega_{\mathcal{X}/\Delta}^k$, and, Δ being smooth of dimension 1, the $\pi_*\Omega_{\mathcal{X}/\Delta}^k$ are in fact locally free. This implies that also $\mathcal{H} := \pi_*d\mathcal{O}_{\mathcal{X}/\Delta}$ is locally free since it is a subsheaf of a locally free sheaf on Δ . Any section in $\mathcal{H}$ is a holomorphic differential form on $\mathcal{X}$ which restricts to a closed form on any fibre.

The base change map

$$\pi_*\Omega_{\mathcal{X}/\Delta}^k \otimes \mathbb{C}_t \hookrightarrow H^0(\mathcal{X}_t, \Omega_{\mathcal{X}/\Delta|_{\mathcal{X}_t}}^k) = H^0(\mathcal{X}_t, \Omega_{\mathcal{X}_t}^k)$$

is an injection, since $\{t\}$ has codimension 1 in Δ ([GR84], Prop. 2, p. 208, p. 209). By possibly shrinking our disk we may assume that on Δ^* the dimensions of $H^0(\mathcal{X}_t, \Omega_{\mathcal{X}_t}^k)$ are constant ($k = 1, 2$) and the map of vector bundles $d_v : \pi_*\Omega_{\mathcal{X}/\Delta}^1 \rightarrow \pi_*\Omega_{\mathcal{X}/\Delta}^2$ has constant rank. Then also $h^0(\mathcal{X}_t, d\mathcal{O}_{\mathcal{X}_t})$ is constant on Δ^* and we have an isomorphism

$$\pi_*d\mathcal{O}_{\mathcal{X}/\Delta} \otimes \mathbb{C}_t \cong H^0(\mathcal{X}_t, d\mathcal{O}_{\mathcal{X}_t})$$

for $t \neq 0$. In particular $\mathcal{H}$ has rank $q = h^0(\mathcal{X}_{t_\nu}, d\mathcal{O}_{\mathcal{X}_{t_\nu}})$.

The map $\mathcal{H} \hookrightarrow H^1(\mathcal{X}_0, \mathbb{C}) \otimes \mathcal{O}_\Delta$ is induced by the inclusion

$$H^0(\mathcal{X}_t, d\mathcal{O}_{\mathcal{X}_t}) \hookrightarrow H^1(\mathcal{X}_0, \mathbb{C})$$

on each fibre. □

Considering a product of elliptic curves deforming to a simple torus it is clear that a very good Albanese-Quotient may not be stable under small deformations. On the other hand we have:

Proposition 2.16 — *Let $\pi : \mathcal{X} \rightarrow \Delta$ be a smooth family of compact, complex manifolds over the unit disk and let $V \subset H_1(\mathcal{X}_0, \mathbb{Q})$. Then the set*

$$Q := \{t \in \Delta \mid V \text{ defines a very good Albanese-Quotient on } \mathcal{X}_t\}$$

is a (possibly empty) analytic subset of Δ .

Proof. The question is local on Δ and hence we may assume that we are in the situation of Lemma 2.15. Consider the composition map of vector bundles on Δ given by

$$\phi : \mathcal{H} \rightarrow H^1(\mathcal{X}_0, \mathbb{C}) \otimes \mathcal{O}_\Delta \rightarrow (H^1(\mathcal{X}_0, \mathbb{C})/Ann(V_{\mathbb{C}})) \otimes \mathcal{O}_\Delta.$$

If $\dim_{\mathbb{C}} Ann(V_{\mathbb{C}}) = 2q$ then V induces a very good Quotient Albanese map on $\mathcal{X}_t$ if and only if the kernel of the map ϕ_t has dimension q which is the maximal possible dimension (see Remark 2.6). Writing ϕ as a matrix with holomorphic entries we see that this is equivalent to the vanishing of the determinants of all minors of a certain dimension which is an analytic condition. □

Corollary 2.17 — *Let $\pi : \mathcal{X} \rightarrow \Delta$ be a smooth family of compact, complex manifolds over the unit disk and let $(t_\nu)_{\nu \in \mathbb{N}}$ be a sequence converging to some point in Δ . If $V \subset H_1(\mathcal{X}_0, \mathbb{Q})$ gives rise to a very good Albanese-Quotient on $\mathcal{X}_t$, for $t = t_\nu$ then this holds for all $t \in \Delta$.*

If the quotient Albanese map is surjective for some t_ν then it is surjective for all t .

Proof. Consider the set Q as in the Proposition. An analytic subset which has an accumulation point must have dimension at least 1. But since Δ itself is 1-dimensional we have $Q = \Delta$ and the first claim is proved. The last statement follows immediately from 2.14. $\square$

We can now study to what extent properties of the general fibre transfer to the central fibre. First we show that there is indeed a family of fibrations.

Lemma 2.18 — *Let $\pi : \mathcal{X} \rightarrow \Delta$ be a smooth family over a small disc and let $V \subset H_1(\mathcal{X}_0, \mathbb{Q})$ be a subspace inducing a very good Albanese-Quotient on every fibre. Then after possibly shrinking Δ there is a family of tori $\pi' : \mathcal{B} \rightarrow \Delta$ and a map Φ inducing a diagram*

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{\Phi} & \mathcal{B} \\ \pi \searrow & & \swarrow \pi' \\ & \Delta & \end{array}$$

such that for every $t \in \Delta$ the map $\Phi_t : \mathcal{X}_t \rightarrow \mathcal{B}_t$ is the quotient Albanese map.

We say that Φ is a family of Albanese-Quotients.

Proof. We may assume that we are in the situation of Lemma 2.15 and that there is a local cross section $s : \Delta \rightarrow \mathcal{X}$. We define $\mathcal{H}' := \text{Ann}(V_{\mathbb{C}}) \cap \mathcal{H}$ and get our family $\mathcal{B} \rightarrow \Delta$ by taking the quotient of $\mathcal{H}'^*$ by the image of $H_1(X, \mathbb{Z}) / (H_1(X, \mathbb{Z}) \cap V)$. The map Φ can be defined by mapping a point $x \in \mathcal{X}$ to the map $\omega \mapsto \int_{\gamma_x} \omega$ where γ_x is any path joining x to $s(\Delta)$. Then Φ restricts to the quotient Albanese map on every fibre as claimed. $\square$

Proposition 2.19 — *Let*

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{\Phi} & \mathcal{B} \\ \pi \searrow & & \swarrow \pi' \\ & \Delta & \end{array}$$

be a family of Albanese-Quotients such that $\Phi_t : \mathcal{X}_t \rightarrow \mathcal{B}_t$ is a smooth holomorphic fibration for $t = t_\nu$ where $(t_\nu)_{\nu \in \mathbb{N}}$ is a sequence converging to 0 in Δ . Then there is a small neighbourhood Δ' of zero such that Φ_t is a smooth holomorphic fibration for all $t \in \Delta'$.

Proof. By Proposition 2.17 the map Φ_0 is also surjective. Recall that Φ is given by integration over the closed holomorphic 1-forms in

$$U := \text{Ann}(V_{\mathbb{C}}) \cap H_0(d\mathcal{O}_{\mathcal{X}}).$$

Let $\omega_1, \dots, \omega_m$ be a basis for U . Then $\omega := \omega_1 \wedge \dots \wedge \omega_m$ generates a sub line bundle of $\Omega_{\mathcal{X}}^m$, namely the pullback $\Phi^* K_{\mathcal{B}/\Delta}$ of the relative canonical bundle of the family $\mathcal{B} \rightarrow \Delta$.

The rank of the Jacobian of Φ in some point $p \in \mathcal{X}$ is not maximal if and only if ω vanishes in p , i.e. in the points of the zero divisor $R := Z(\omega)$. But since Φ_t is smooth for $t = t_\nu$ the divisor R is completely contained in a union of fibres and we can choose Δ' such that Φ_t is smooth for $t \in \Delta' \setminus \{0\}$.

Hence we may assume that R is completely contained in $\mathcal{X}_0$ and, since $\mathcal{X}_0$ is irreducible of codimension 1, we have in fact $R = \mathcal{X}_0$ or $R = \emptyset$.

If R is not empty then there is a maximal $k \in \mathbb{N}$ such that ω/t^k is holomorphic and hence, after a base change $\Delta \xrightarrow{t^k} \Delta$, there is at least one point in $\mathcal{X}_0$ where ω does not vanish. But then it can vanish nowhere by dimension reasons and this proves that the Jacobian has maximal rank everywhere and the central fibre is indeed a smooth holomorphic fibration. $\square$

Now we want to know, when fibration on the central fibre is locally free provided this holds for Φ_t ($t \neq 0$). For a family of manifolds $\Phi : \mathcal{X} \rightarrow \mathcal{B}$ we define the sheaf of relative tangent vectors $\Theta_{\mathcal{X}/\mathcal{B}}$ by the sequence

$$0 \rightarrow \Theta_{\mathcal{X}/\mathcal{B}} \rightarrow \Theta_{\mathcal{X}} \rightarrow \Phi^* \Theta_{\mathcal{B}} \rightarrow 0.$$

Proposition 2.20 — *Let*

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{\Phi} & \mathcal{B} \\ & \searrow \pi \quad \swarrow \pi' & \\ & \Delta & \end{array}$$

be a family of smooth holomorphic fibrations over tori, parametrised by the unit disc, such that all the fibres of the maps $\Phi_t : \mathcal{X}_t \rightarrow \mathcal{B}_t$ are contained in a class $\mathcal{C}$ of compact complex manifolds.

If Φ_t is locally trivial for $t \neq 0$ then also Φ_0 is locally trivial if one of the following conditions holds:

- (i) *There is a coarse moduli space $\mathfrak{M}_{\mathcal{C}}$ for manifolds in class $\mathcal{C}$ which is Hausdorff.*
- (ii) *For every manifold in $Y \in \mathcal{C}$ there is a local moduli space, in other words, the Kuranishi family of Y is universal.*
- (iii) *The sheaf $R^1 \Phi_* \Theta_{\mathcal{X}/\mathcal{B}}$ is locally free. (This holds if $h^1(\Phi^{-1}(p), \Theta_{\mathcal{X}_p})$ does not depend on $p \in \mathcal{B}$.)*

Proof. In the first case let F, F' be two fibres of Φ_0 . By [FG65] it suffices to show that $F \cong F'$. By pullback we can obtain two families $\mathcal{F}, \mathcal{F}'$ of manifolds in $\mathcal{C}$ parametrised by Δ such that $\mathcal{F}_t$ is a fibre of Φ_t and with central fibres F and F' respectively. Since Φ_t is locally trivial for $t \neq 0$ by assumption the corresponding moduli maps to $\mathfrak{M}_{\mathcal{C}}$ coincide for $t \neq 0$ and since $\mathfrak{M}_{\mathcal{C}}$ was assumed to be Hausdorff they have to coincide also for $t = 0$ thence $F \cong F'$ as claimed.

For the second case we consider the family $\mathcal{X} \rightarrow \mathcal{B}$ in the neighbourhood of some point $p_0 \in \mathcal{B}_0$.

Let $\text{Kur}(\Phi^{-1}(p_0))$ be the Kuranishi family of $\Phi^{-1}(p_0)$ which is universal by assumption, i.e., in some neighbourhood U of p_0 we get a unique moduli map $\mu : U \rightarrow \text{Kur}(\Phi^{-1}(p_0))$ such that the point $\mu(p)$ corresponds to the manifold $\Phi^{-1}(p)$.

If we choose U sufficiently small we can find local coordinates $x = (t, y)$ around p_0 such that $\pi'(x) = t$. Since $\Phi_t : \mathcal{X}_t \rightarrow \mathcal{B}_t$ is locally trivial for $t \neq 0$ we have

$$\frac{\partial \mu(t, y)}{\partial y} = 0$$

on the dense set where $t \neq 0$ and hence everywhere.

Therefore the moduli map is constant on $U \cap \mathcal{B}_0$ and, since the point $p_0 \in \mathcal{B}_0$ was arbitrary and $\mathcal{B}_0$ is connected, all the fibres of ϕ_0 are isomorphic and Φ_0 is locally trivial by [FG65].

It remains to treat the last case. The fibration $\Phi_0 : \mathcal{X}_0 \rightarrow \mathcal{B}_0$ is locally trivial if and only if the Kodaira-Spencer map

$$\rho_0 : \Theta_{\mathcal{B}_0} \rightarrow R^1\Phi_{0*}\Theta_{\mathcal{X}_0/\mathcal{B}_0}$$

vanishes identically. ([GR84], Proposition 1, p. 208).

We want to study the relation between ρ_0 and the Kodaira-Spencer map ρ for the whole family via the base change homomorphism. In particular we are interested in the subsheaf $\Theta_{\mathcal{B}/\Delta} \subset \Theta_{\mathcal{B}}$ that restricts to $\Theta_{\mathcal{B}_t}$ on every fibre of π' .

Let

$$\rho' : \Theta_{\mathcal{B}/\Delta} \rightarrow \Theta_{\mathcal{B}} \xrightarrow{\rho} R^1\Phi_*\Theta_{\mathcal{X}/\mathcal{B}}$$

be the composition map which is a map of vector bundles by our assumptions. We claim that ρ' is in fact identically zero:

Let Z be an analytic subspace of $\mathcal{B}$ of codimension 1 and let $\mathcal{I}$ be the corresponding ideal sheaf. Then for any sheaf $\mathcal{G}$ on $\mathcal{X}$ there is the base change map

$$\Phi_{\mathcal{I}} : R^1\Phi_*\mathcal{G}/\mathcal{I} \cdot R^1\Phi_*\mathcal{G} \rightarrow R^1\Phi_*(\mathcal{G}|_{\Phi^{-1}Z})$$

which is injective ([GR84], Prop. 2, p. 208, p. 209).

For our subspaces $\mathcal{B}_t$ the naturality of the base change map yields a commutative diagram

$$\begin{array}{ccc} \Theta_{\mathcal{B}/\Delta}|_{\mathcal{B}_t} & \xrightarrow{\rho'} & (R^1\Phi_*\Theta_{\mathcal{X}/\mathcal{B}})|_{\mathcal{B}_t} \\ \downarrow \cong & & \downarrow \\ \Theta_{\mathcal{B}_t} & \xrightarrow{\rho_t} & R^1\Phi_{t*}(\Theta_{\mathcal{X}_t/\mathcal{B}_t}). \end{array}$$

If $t \neq 0$ then we have $\rho_t \equiv 0$ since Φ_t is locally trivial. Therefore the map of vector bundles $\rho' \equiv 0$ because it vanishes on the dense open set $\mathcal{B} \setminus \mathcal{B}_0$.

Looking again at the diagram for $t = 0$ we see that also ρ_0 must be zero. Hence Φ_0 is a locally trivial fibration as claimed. $\square$

Remark 2.21 — By a theorem of Wavrik [Wav69] condition (ii) in Proposition 2.20 holds if $h^k(\Phi^{-1}(p), \Theta_{\mathcal{X}_p})$ does not depend on $p \in \mathcal{B}$ for $k = 0$ or $k = 1$ and the Kuranishi space of $\Phi^{-1}(p)$ is reduced for all $p \in \mathcal{B}$.

Proof of Theorem 2.11. Let $\mathcal{C}$ be a good fibre class and $\pi : \mathcal{X} \rightarrow \Delta$ be a family of compact, complex manifolds. Let $(t_\nu)_{\nu \in \mathbb{N}}$ be a sequence in Δ converging to zero such that subspace $V \subsetneq H_1(X, \mathbb{Q})$ is $\mathcal{C}$ -fibering on $\mathcal{X}_{t_\nu}$ for all ν .

In particular V defines a very good Albanese-Quotient on $\mathcal{X}_t$ for $t = t_\nu$ and hence for all $t \in \Delta$ by Corollary 2.17. So we may assume that we are in the situation of Lemma 2.18.

Our assumptions guarantee that we can first apply Proposition 2.19 and then Proposition 2.20. This concludes the proof of the theorem. $\square$

3. COMPLEX STRUCTURES ON CERTAIN LIE-ALGEBRAS

The aim of this section is to study the possible complex structures on certain types of Lie-algebras. We are particularly interested in the existence of SPTBS in Lie-algebras in view of Corollary 1.12 and Theorem 2.12.

Using only the dimensions of the subspaces in the descending and ascending central series we will try to give a complete picture for Lie-algebras with commutator subalgebra of dimension at most three. By giving lots of examples we will also show that our classification cannot be improved without considering other properties of the Lie-algebras. In the case $\dim \mathcal{C}^1 \mathfrak{g} = 1$ we can show that there is a unique complex structure up to isomorphism.

3.1. Notations and basic results. In the sequel $\mathfrak{g}$ will denote a nilpotent Lie-algebra and J a complex structure on $\mathfrak{g}$ which is always assumed to be integrable in the sense of Definition 1.1; in particular $\dim \mathfrak{g}$ is always even.

We continue to use the notation introduced in Section 1, see Section 1.4 for the notation in the examples.

For later reference we collect some basic facts about complex structures and Lie-algebras in the next lemma.

Lemma 3.1 — *Let $(\mathfrak{g}, J)$ be a real Lie-algebra with a complex structure.*

- (i) *Let $V \subset \mathfrak{g}$ be a real subspace. Then the following are equivalent:*
 - *V is J -invariant.*
 - *V is a complex subspace of $(\mathfrak{g}, J)$ considered as a complex vector space.*
 - *$V_{\mathbb{C}} = (V_{\mathbb{C}} \cap \mathfrak{g}^{1,0}) \oplus (V_{\mathbb{C}} \cap \mathfrak{g}^{0,1})$.*
- (ii) *If $V \subset \mathfrak{g}$ is J -invariant then $\dim_{\mathbb{R}} V$ is even. In particular if $V \subset \mathfrak{g}$ is a nontrivial, J -invariant subspace contained in a real 2-dimensional subspace W then $V = W$.*
- (iii) *Let V be a real subspace of $\mathfrak{g}$. Then the largest J -invariant subspace of V is*

$$((V_{\mathbb{C}} \cap \mathfrak{g}^{1,0}) \oplus (V_{\mathbb{C}} \cap \mathfrak{g}^{0,1})) \cap \mathfrak{g}$$

where $\mathfrak{g}$ is identified with $\{x \otimes 1 \mid x \in \mathfrak{g}\} \subset \mathfrak{g}_{\mathbb{C}}$.

- (iv) *For any $x \in \mathfrak{g}$ holds: $\text{ad}_x(-) = [x, -] \neq 0 \Leftrightarrow x \notin \mathcal{Z}\mathfrak{g}$.*
- (v) *If $\mathfrak{g}$ is 2-step nilpotent then $\mathcal{C}^1 \mathfrak{g} \subset \mathcal{Z}\mathfrak{g}$.*
- (vi) *Let $V \subset \mathfrak{g}$ be a real subspace with $\dim_{\mathbb{R}} V = 3$. If $W_1, W_2 \subset V$ are both non-trivial, J -invariant subspaces, then $W_1 = W_2$.*
- (vii) *The complex structure is abelian, i.e., $\mathfrak{g}^{1,0}$ is an abelian subalgebra of $\mathfrak{g}_{\mathbb{C}}$, if and only if $[x, y] = [Jx, Jy]$ for all $x, y \in \mathfrak{g}$. In this case the ascending central series $(\mathcal{Z}^i \mathfrak{g})$ is J -invariant.*
- (viii) *If $\mathcal{C}^1 \mathfrak{g}$ contains no J -invariant subspace then the complex structure J is abelian.*

Proof. We prove only the last three assertions, the rest being clear.

- (vi) Both W_1 and W_2 have positive, even real dimension, hence

$$\dim_{\mathbb{R}} W_1 = \dim_{\mathbb{R}} W_2 = 2 \text{ and } W_1 \cap W_2 \neq 0$$

by dimensional reasons. Since the intersection $W_1 \cap W_2$ is again J -invariant it has also dimension 2 by (ii) and we have

$$W_1 = W_1 \cap W_2 = W_2$$

as claimed.

(vii) This is a straightforward calculation which can be found for example in [MPPS06].

(viii) Since we assumed J to be integrable $\mathfrak{g}^{1,0}$ is a subalgebra of $\mathfrak{g}_{\mathbb{C}}$ and hence

$$\mathcal{C}^1 \mathfrak{g}^{1,0} \subset \mathcal{C}^1 \mathfrak{g}_{\mathbb{C}} \cap \mathfrak{g}^{1,0} \subset \mathcal{C}^1 \mathfrak{g}_{\mathbb{C}}.$$

If $\mathcal{C}^1 \mathfrak{g}$ contains no J -invariant subspace then $\mathcal{C}^1 \mathfrak{g}_{\mathbb{C}} \cap \mathfrak{g}^{1,0} = 0$ by (i). $\square$

The main tool in our analysis will be the following technical lemma.

Lemma 3.2 — *Let $\mathfrak{g}$ be a nilpotent Lie-algebra with complex structure J . Let $\mathcal{Z}_J \mathfrak{g} := \mathcal{Z} \mathfrak{g} + J \mathcal{Z} \mathfrak{g}$ be the smallest complex subspace of $\mathfrak{g}$ which contains the centre, $\mathcal{W}_i := \mathcal{Z}_J \mathfrak{g} \cap \mathcal{C}^i \mathfrak{g}$ and*

$$\mathcal{V}_{i+1}^J := [\mathcal{W}_i, \mathfrak{g}] = \text{span}_{\mathbb{R}}\{[w, x] \mid w \in \mathcal{W}_i, x \in \mathfrak{g}\}.$$

Then:

- (i) $\mathcal{V}_i^J$ is a (possibly trivial) J -invariant subspace of the i -th commutator subalgebra $\mathcal{C}^i \mathfrak{g}$ and we have $\mathcal{V}_{i+1}^J \subset \mathcal{V}_i^J$.
- (ii) If there is some $x \in \mathcal{W}_i \setminus \mathcal{Z} \mathfrak{g}$ then $\mathcal{V}_{i+1}^J \neq 0$.
- (iii) The centre is J -invariant if and only if $\mathcal{V}^J := \mathcal{V}_1^J = 0$.

If the complex structure is fixed we will often omit it from the notation.

Proof. We begin with the last assertion: the centre is not fixed by J if and only if for some $z \in \mathcal{Z} \mathfrak{g}$ the element Jz is not in the centre, which means that there exists some $x \in \mathfrak{g}$ such that $[Jz, x] \neq 0$ which is equivalent to $\mathcal{V}^J \neq 0$.

For (i) we only have to show that $x \in \mathcal{V}_i^J$ implies $Jx \in \mathcal{V}_i^J$. We will do this on generators of the form $x = [Jz, y]$ for some z in the centre of $\mathfrak{g}$ and $y \in \mathfrak{g}$. The Nijenhuis tensor then implies

$$Jx = J[Jz, y] = [Jz, Jy] - [z, y] - J[z, Jy] = [Jz, Jy] \in \mathcal{V}_i^J.$$

The second assertion follows immediately from the definition of $\mathcal{V}_i^J$ and 3.1 (iv). $\square$

In some cases the lemma will enable us to prove that there do not exist complex structures on a certain class of Lie-algebras with the following argument: we assume the existence of a complex structure and then deduce that some odd-dimensional subspace should be invariant under J which is impossible.

We give two applications:

Proposition 3.3 — *Let $\mathfrak{g}$ be a 2-step nilpotent Lie-algebra. Then every integrable complex structure on $\mathfrak{g}$ is nilpotent.*

Proof. Let J be a complex structure on $\mathfrak{g}$. We will prove the claim by induction on the dimension of $\mathfrak{g}$. We have to show that for some k we have $\mathcal{T}^k \mathfrak{g} = \mathfrak{g}$ in the minimal torus bundle series or, equivalently, that $\mathfrak{g}/\mathcal{T}^{k-1} \mathfrak{g}$ is an abelian Lie-algebra.

Since $\mathfrak{g}$ is 2-step nilpotent we have

$$\mathfrak{g} \supset \mathcal{Z} \mathfrak{g} \supset \mathcal{C}^1 \mathfrak{g} \supset 0$$

and, by Lemma 3.2, either $\mathcal{V}_1$ or $\mathcal{Z}\mathfrak{g}$ is a nontrivial J -invariant subspace of the centre. In particular the centre has dimension at least 2 and if $\dim \mathcal{Z}\mathfrak{g} = 2$ then the centre is J -invariant (3.1 (ii)). This proves the claim if $\mathfrak{g}$ has dimension 4 since the quotient is then necessarily abelian.

In higher dimension we have $\mathcal{T}^1\mathfrak{g} \neq 0$. If $\mathfrak{g}/\mathcal{T}^1\mathfrak{g}$ is abelian we are done, else $\mathfrak{g}/\mathcal{T}^1\mathfrak{g}$ is still 2-step nilpotent and we can use the induction hypothesis. $\square$

Proposition 3.4 — *Let $\mathfrak{g}$ be a nilpotent Lie-algebra and $k = \dim \mathcal{Z}\mathfrak{g}$. Let $m = k + 1$ if k is odd and $m = k + 2$ if k is even. Assume that $\mathcal{Z}\mathfrak{g} = \mathcal{C}^1\mathfrak{g}$ and that one of the following holds:*

- (i) *For every m -dimensional subspace W , which contains the centre, we have $[W, \mathfrak{g}] = \mathcal{C}^1\mathfrak{g}$.*
- (ii) *The map $\text{ad}_x : \mathfrak{g} \rightarrow \mathcal{C}^1\mathfrak{g}$ is surjective for all $x \notin \mathcal{Z}\mathfrak{g}$.*

If $k = \dim(\mathcal{Z}\mathfrak{g})$ is not even then there does not exist any integrable complex structure on $\mathfrak{g}$ and if $k = \dim(\mathcal{Z}\mathfrak{g})$ is even then any complex structure on such a $\mathfrak{g}$ is nilpotent and

$$\mathfrak{g} \supset \mathcal{Z}\mathfrak{g} = \mathcal{C}^1\mathfrak{g} \supset 0$$

is a SPTBS for $\mathfrak{g}$.

Proof. Clearly the (ii) implies the (i). Now assume we have a complex structure J on $\mathfrak{g}$ such that the centre is not J -invariant. Then $\mathcal{Z}_J\mathfrak{g} := \mathcal{Z}\mathfrak{g} + J\mathcal{Z}\mathfrak{g}$ is an even dimensional subspace such that $\mathcal{Z}\mathfrak{g} \subsetneq \mathcal{Z}_J\mathfrak{g}$ and therefore it has dimension at least m . The subspace $\mathcal{V}^J$ is nonempty and by assumption (i) it is in fact equal to $\mathcal{C}^1\mathfrak{g} = \mathcal{Z}\mathfrak{g}$. But then Lemma 3.2 implies that $\mathcal{Z}\mathfrak{g} = \mathcal{V}^J$ is J -invariant – a contradiction.

It follows that the centre is J -invariant for every complex structure J on $\mathfrak{g}$. Therefore the centre cannot have odd dimension if there exists a complex structure. $\square$

In Section 4.2 we will also need the following:

Lemma 3.5 — *Let $(\mathfrak{g}, J)$ be a nilpotent Lie-algebra with a complex structure, $z \in \mathcal{Z}\mathfrak{g}$ and $W := \langle z, Jz \rangle$. Then $(\text{im } \text{ad}_{Jz}) \cap W = 0$.*

Proof. Since $\mathfrak{g}$ is nilpotent $Jz \notin \text{im } \text{ad}_{Jz}$ and it remains to prove that $z \notin \text{im } \text{ad}_{Jz}$. So assume that there is some $x \in \mathfrak{g}$ such that $[Jz, x] = z$. Then with the same calculation as in Lemma 3.2 we have

$$\text{ad}_{Jz}(Jx) = [Jz, Jx] = J[Jz, x] = Jz,$$

so $Jz \in \text{im } \text{ad}_{Jz}$ which is a contradiction. $\square$

3.2. The case $\dim(\mathcal{C}^1\mathfrak{g}) = 1$. Recall that a Heisenberg algebra H_{2n+1} is a nilpotent Lie-algebra which admits a basis $x_1, \dots, x_n, y_1, \dots, y_n, c$ such that c is central and the structure equations are

$$(3) \quad \begin{aligned} [x_i, y_i] &= -[y_i, x_i] = c, & i &= 1, \dots, n, \\ [x_i, x_j] &= -[y_i, y_j] = 0, & i, j &= 1, \dots, n. \end{aligned}$$

Proposition 3.6 — *Let $\mathfrak{g}$ be a real nilpotent Lie-algebra with $\dim(\mathcal{C}^1\mathfrak{g}) = 1$. Then $\mathfrak{g}$ is 2-step nilpotent and $\mathfrak{g} \cong H_{2n+1} \oplus \mathbb{R}^m$ is the direct sum of a Heisenberg algebra with an abelian Lie-algebra with generators $z_1, \dots, z_m$. Then for any choice of signs*

$$(4) \quad \begin{aligned} Jx_i &:= \pm y_i & (i = 1, \dots, n), \\ Jz_{2k-1} &:= z_{2k} & (k = 1, \dots, r), \\ Jc &:= z_{2r+1}, & J^2 = -id_{\mathfrak{g}}, \end{aligned}$$

defines a complex structure on $\mathfrak{g}$ and every complex structure on $\mathfrak{g}$ is equivalent to a complex structure of this kind. Thus, every complex structure is abelian and the ascending central series is a SPTBS for $\mathfrak{g}$.

The centre has necessarily even dimension since we assumed $\mathfrak{g}$ to have even dimension.

Proof. Let $c \in \mathcal{C}^1\mathfrak{g}$ be a generator of the commutator subalgebra. Considering the Lie-bracket as an alternating form on $\mathfrak{g}/\langle c \rangle$ the classification of alternating forms on vector spaces. ([Lan84], XIV, 9) yields the claimed decomposition.

Now assume that $\mathfrak{g}$ admits an integrable complex structure J . The commutator $\mathcal{C}^1\mathfrak{g}$ cannot contain any complex subspace for dimensional reasons and writing the Nijenhuis tensor as

$$[Jx, Jy] - [x, y] = J([Jx, y] + [x, Jy])$$

we see that the left hand side is in $\mathcal{C}^1\mathfrak{g}$ while the right hand side is not. This yields

$$[Jx, Jy] = [x, y]$$

for all $x, y \in \mathfrak{g}$ and hence any complex structure on $\mathfrak{g}$ is abelian by Lemma 3.1 (vii).

It is a straightforward calculation to show that J as in (4) defines a complex structure and it remains to show that every integrable complex structure can be written in this way with respect to a suitable basis.

Assume that we have $\mathfrak{g} \cong H_{2n+1} \oplus \mathbb{R}^{2r+1}$ as above and we are given a complex structure J on $\mathfrak{g}$. The centre $\mathcal{Z}\mathfrak{g}$ is J -invariant and we can choose a basis for the centre such that

$$Jz_{2k-1} = z_{2k} \quad (k = 1, \dots, r), \quad Jc = z_{2r+1}.$$

Let us fix an arbitrary complex subspace V such that $\mathfrak{g} = V \oplus \mathcal{Z}\mathfrak{g}$. The remaining elements of the basis are provided by the following:

Claim: If (V, J) is a real vector space with a complex structure and

$$[-, -] : V \times V \rightarrow \mathbb{R} \cdot c$$

a non-degenerate alternating bilinear form on V , such that $[x, y] = [Jx, Jy]$ for all $x, y \in V$ then there exists a basis x_i, y_i of V which satisfies (3) and (4).

We will prove our claim by induction on the dimension of V . First of all we show that we can always find an $x \in V$ such that $[x, Jx] \neq 0$.

We pick a nonzero $a \in V$. If $[a, Ja] \neq 0$ then we are done and hence assume $[a, Ja] = 0$. There is some $b \in V$ such that $[a, b] = c$ since the bracket is non-degenerate. If $[b, Jb] \neq 0$ we set $x = b$.

Otherwise also $[b, Jb] = 0$ and we calculate using $[Ja, Jb] = [a, b]$

$$\begin{aligned} [a + Jb, J(a + Jb)] &= [a + Jb, Ja - b] \\ &= [a, Ja] - [a, b] + [Jb, Ja] - [Jb, b] \\ &= -2[a, b] = -2c \neq 0, \end{aligned}$$

i.e., we can set $x = a + Jb$.

If $[x, Jx] \neq 0$ then $[x, Jx] = \lambda c$ for some $\lambda \in \mathbb{R} \setminus \{0\}$. Setting $x_1 = x/\sqrt{|\lambda|}$ and $y_1 := \text{sign}(\lambda)Jx_1$ (where $\text{sign}(\lambda)$ is the sign of λ) we have

$$[x_1, y_1] = c \text{ and } Jx_1 = \text{sign}(\lambda)y_1.$$

If $\dim_{\mathbb{R}} V = 2$ we are done; otherwise we can apply the induction hypothesis to the subspace orthogonal to $\langle x_1, y_1 \rangle$ with respect to the Lie-bracket. This concludes the proof. $\square$

3.3. The case $\dim(\mathcal{C}^1 \mathfrak{g}) = 2$. We begin with a preliminary lemma.

Lemma 3.7 — *Let $\mathfrak{g}$ be a nilpotent Lie-algebra with $\dim(\mathcal{C}^1 \mathfrak{g}) = 2$ and J a complex structure on $\mathfrak{g}$. Then at least one of the subspaces $\mathcal{Z}\mathfrak{g}$ and $\mathcal{C}^1 \mathfrak{g}$ is J -invariant.*

Proof. We have $0 \subset \mathcal{V} \subset \mathcal{C}^1 \mathfrak{g}$ and since $\mathcal{V}$ is even dimensional and $\dim(\mathcal{C}^1 \mathfrak{g}) = 2$, we have either $\mathcal{V} = 0$ or $\mathcal{V} = \mathcal{C}^1 \mathfrak{g}$. An application of Lemma 3.2 (iii) concludes the proof. $\square$

The classification in this case reads as follows.

Proposition 3.8 — *Let $\mathfrak{g}$ be a nilpotent Lie-algebra with $\dim(\mathcal{C}^1 \mathfrak{g}) = 2$. Then every complex structure on $\mathfrak{g}$ is nilpotent and we have the following cases.*

- (i) *If $\mathfrak{g}$ is 3-step nilpotent then the following holds: if one of the subalgebras $\mathcal{Z}^i \mathfrak{g}$ has odd dimension, then $\mathfrak{g}$ does not admit any complex structure. If there exists a complex structure J on $\mathfrak{g}$ then J is abelian and the ascending central series is a SPTBS.*
- (ii) *If $\mathfrak{g}$ is 2-step nilpotent and $\dim(\mathcal{Z}\mathfrak{g})$ is odd then*

$$0 \subset \mathcal{C}^1 \mathfrak{g} \subset \mathfrak{g}$$

is a SPTBS on $\mathfrak{g}$.

- (iii) *If $\mathfrak{g}$ is 2-step nilpotent and $\dim(\mathcal{Z}\mathfrak{g})$ is even then for every complex structure on $\mathfrak{g}$ either*

$$0 \subset \mathcal{C}^1 \mathfrak{g} \subset \mathfrak{g} \quad \text{or} \quad 0 \subset \mathcal{Z}\mathfrak{g} \subset \mathfrak{g}$$

is a torus bundle series on $\mathfrak{g}$ but a SPTBS does not exist in general if $\dim(\mathcal{Z}\mathfrak{g}) \geq 4$. If $\dim(\mathcal{Z}\mathfrak{g}) = 2$ the two series coincide and we have a SPTBS.

Proof. We treat the cases separately:

- (i) Assume that we have a complex structure J on $\mathfrak{g}$ and $\mathfrak{g}$ is 3-step nilpotent. It suffices to show that $\mathcal{C}^1 \mathfrak{g}$ is not a J -invariant subspace by Lemma 3.1 (vii), (viii) since $\dim_{\mathbb{R}} \mathcal{C}^1 \mathfrak{g} = 2$.

Assume the contrary. Writing $\mathcal{C}^2 \mathfrak{g} = \langle c_2 \rangle$ we then have $\mathcal{C}^1 = \langle c_1 := Jc_2, c_2 \rangle$, in particular $\mathcal{W}_1 \neq 0$ in the notation of Lemma 3.2.

But $c_2 \in \text{im}(ad_{c_1})$ and hence, by Lemma 3.2 (ii), $\mathcal{V}_2 \neq 0$ which is impossible since $\dim \mathcal{C}^2 \mathfrak{g} = 1$.

- (ii) Clearly $\mathcal{Z}\mathfrak{g}$ can never be J -invariant for any complex structure on $\mathfrak{g}$ if $\dim \mathcal{Z}\mathfrak{g}$ is odd and hence the assertion follows from Lemma 3.7.
- (iii) We observe that 2-step nilpotency implies $\mathcal{C}^1 \mathfrak{g} \subset \mathcal{Z}\mathfrak{g}$ and hence both subspaces coincide if $\dim \mathcal{C}^1 \mathfrak{g} = \dim \mathcal{Z}\mathfrak{g} = 2$. So by Lemma 3.7

$$0 \subset \mathcal{C}^1 \mathfrak{g} = \mathcal{Z}\mathfrak{g} \subset \mathfrak{g}$$

is a SPTBS in this case. The remaining assertions follow from Example 3.9 given below.

□

Example 3.9 — Consider the following 10-dimensional Lie-algebra given by a basis $\mathfrak{g} = \langle e_1, e_2, e_3, f_1, f_2, f_3, z_1, z_2, c_1, c_2 \rangle$ and the structure equations

$$dc^1 = e^{12} + f^{12}, \quad dc^2 = e^{13} + f^{13}$$

with respect to the dual basis. $\mathfrak{g}$ is a 2-step nilpotent Lie-algebra with centre $\mathcal{Z}\mathfrak{g} = \langle z_1, z_2, c_1, c_2 \rangle$ and commutator $\mathcal{C}^1 \mathfrak{g} = \langle c_1, c_2 \rangle$. We give three interable complex structures J_1, J_2 and J_3 on $\mathfrak{g}$ by

$$\begin{array}{lll} J_1 e_1 = f_1, & J_2 z_1 = e_1, & J_3 e_1 = f_1, \\ J_1 e_2 = f_2, & J_2 z_2 = f_1, & J_3 e_2 = f_2, \\ J_1 e_3 = f_3, & J_2 c_1 = c_2, & J_3 e_3 = f_3, \\ J_1 z_1 = c_1, & J_2 e_2 = e_3, & J_3 z_1 = z_2, \\ J_1 z_2 = c_2, & J_2 f_2 = f_3, & J_3 c_1 = c_2. \end{array}$$

The complex structure J_1 leaves the commutator invariant but not the centre ($\mathcal{V} = \mathcal{C}^1 \mathfrak{g}$), J_2 the centre but not the commutator while J_3 leaves both subspaces invariant. This realises all possible combinations in Proposition 3.8 (iii).

3.4. The case $\dim(\mathcal{C}^1 \mathfrak{g}) = 3$.

Proposition 3.10 — Let $\mathfrak{g}$ be a nilpotent Lie-algebra with $\dim(\mathcal{C}^1 \mathfrak{g}) = 3$. Then the following cases can occur:

- (i) If $\mathfrak{g}$ is 4-step nilpotent then the following holds: if one of the subalgebras $\mathcal{Z}^i \mathfrak{g}$ has odd dimension, then $\mathfrak{g}$ does not admit any complex structure. If there exists a complex structure J on $\mathfrak{g}$ then J is abelian and the ascending central series is a SPTBS.
- (ii) If $\mathfrak{g}$ is 3-step nilpotent and $\dim(\mathcal{C}^2 \mathfrak{g}) = \dim(\mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z}\mathfrak{g}) = 1$ then for any complex structure J on $\mathfrak{g}$ we have $\mathcal{C}^2 \mathfrak{g} \cap \mathcal{V}^J = 0$ but $\mathcal{V}^J \neq 0$ is possible and there may be nilpotent and non-nilpotent complex structures on the same Lie-algebra.
- (iii) If $\mathfrak{g}$ is 3-step nilpotent such that $\dim(\mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z}\mathfrak{g}) = 2$ then:
 - (a) If $\dim \mathcal{Z}^i \mathfrak{g}$ is even for all i then every complex structure J on $\mathfrak{g}$ is nilpotent and one of the subspaces $\mathcal{Z}\mathfrak{g}$ and $\mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z}\mathfrak{g}$ is J -invariant but there is no SPTBS in general.

(b) If $\dim \mathcal{Z}^1 \mathfrak{g}$ is odd or equal to 2 and $\dim \mathcal{Z}^2 \mathfrak{g}$ is even then

$$\mathfrak{g} \supset \mathcal{Z}^2 \mathfrak{g} \supset \mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g} \supset 0$$

is a SPTBS for $\mathfrak{g}$.

(c) If $\dim \mathcal{Z}^2 \mathfrak{g}$ is odd there does not exist any complex structure on $\mathfrak{g}$.

(iv) If $\mathfrak{g}$ is 2-step nilpotent then every complex structure on $\mathfrak{g}$ is nilpotent but in general there does not exist a SPTBS.

Proof. The last assertion follows immediately from Proposition 3.3 and Example 1.14. The remaining cases (i), (ii) and (iii) will be treated separately in the following lemmata. $\square$

Lemma 3.11 — *Let $\mathfrak{g}$ be a 4-step nilpotent Lie-algebra with $\dim(\mathcal{C}^1 \mathfrak{g}) = 3$. If one of the subalgebras $\mathcal{Z}^i \mathfrak{g}$ has odd dimension, then $\mathfrak{g}$ does not admit any complex structure. If there exists a complex structure J on $\mathfrak{g}$ then J is abelian and the ascending central series is a SPTBS.*

Proof. Assume that $\mathfrak{g}$ admits a complex structure J . It suffices to show that $\mathcal{C}^1 \mathfrak{g}$ does not contain a J -invariant subspace by Lemma 3.1 (vii), (viii).

Assume the contrary, i.e., that there is a non-trivial, J -invariant subspace $W \subset \mathcal{C}^1 \mathfrak{g}$ with (necessarily) $\dim_{\mathbb{R}}(W) = 2$. By Lemma 3.1 (vi) every complex subspace of $\mathcal{C}^1 \mathfrak{g}$ is contained in W and equal to W if it is non-trivial. Since $\dim \mathcal{C}^1 \mathfrak{g} = 3$ and $\mathfrak{g}$ is 4-step nilpotent we have

$$\begin{aligned} \dim \mathcal{C}^3 \mathfrak{g} &= 1, & \dim \mathcal{C}^2 \mathfrak{g} &= 2, \\ \mathcal{Z} \mathfrak{g} \cap \mathcal{C}^1 \mathfrak{g} &= \mathcal{C}^3 \mathfrak{g}. \end{aligned}$$

We will now derive a contradiction.

Case 1: $W \cap \mathcal{C}^3 \mathfrak{g} = \langle c_3 \rangle \neq 0$: Then $Jc_3 \in \mathcal{C}^1 \mathfrak{g} \setminus \mathcal{Z} \mathfrak{g}$ and $\mathcal{V}_2 \neq 0$ by Lemma 3.2 (ii). But this in turn implies $\mathcal{C}^2 \mathfrak{g} = \mathcal{V}_2 = W$ for dimensional reasons. Hence $Jc_3 \in \mathcal{C}^2 \mathfrak{g} \setminus \mathcal{Z} \mathfrak{g}$ and again by Lemma 3.2 we have $0 \neq \mathcal{V}_3 \subset \mathcal{C}^3 \mathfrak{g}$ which is impossible since $\mathcal{V}_3$ has dimension at least 2.

Case 2: $W \cap \mathcal{C}^3 \mathfrak{g} = 0$: We construct a basis of $\mathcal{C}^1 \mathfrak{g}$ in the following way: for dimensional reasons W and $\mathcal{C}^2 \mathfrak{g}$ intersect non-trivially and we can choose a nonzero $c_2 \in W \cap \mathcal{C}^2 \mathfrak{g}$. Let $c_1 := Jc_2$ and $c_3 \in \mathcal{C}^3 \mathfrak{g}$ be a generator. Then by our assumptions c_1, c_2, c_3 is a basis of $\mathcal{C}^1 \mathfrak{g}$.

There is $f_2 \in \mathfrak{g}$ such that $[c_2, f_2] = c_3$. The Nijenhuis tensor together with $Jc_2 = -c_1$ yields

$$J(\underbrace{[Jc_2, f_2] + [c_2, Jf_2]}_{\in \mathcal{C}^2 \mathfrak{g}}) = [Jc_2, Jf_2] - [c_2, f_2] \in \mathcal{C}^2 \mathfrak{g}$$

and, since $\mathcal{C}^2 \mathfrak{g} \cap J\mathcal{C}^2 \mathfrak{g} = 0$ under our conditions, both sides are equal to zero. In particular

$$(5) \quad [c_1, f_2] = [Jc_2, f_2] = -[c_2, Jf_2] \in \mathcal{C}^3 \mathfrak{g}$$

is central. Writing $c_1 = [a, b]$ we also have

$$(6) \quad [c_1, c_2] = [[a, b], c_2] = [a, [b, c_2]] - [b, [a, c_2]] \in \mathcal{C}^4 \mathfrak{g} = 0.$$

Since $c_2 \in \mathcal{C}^2\mathfrak{g} \setminus \mathcal{Z}\mathfrak{g}$ we can find $f_1 \in \mathfrak{g}$ and $\lambda \in \mathbb{R}$ such that

$$(7) \quad [c_1, f_1] = c_2 + \lambda c_3.$$

Writing $[f_1, f_2] = \sum_{i=1}^3 \lambda_i c_i$ and using the Jacobi identity we get

$$\begin{aligned} c_3 &= [c_2, f_2] = [c_2 + \lambda c_3, f_2] \\ &\stackrel{(7)}{=} [[c_1, f_1], f_2] \\ &= [c_1, [f_1, f_2]] - [f_1, [c_1, f_2]] \\ &\stackrel{(5)}{=} [c_1, [f_1, f_2]] \\ &= [c_1, \sum_{i=1}^3 \lambda_i c_i] \\ &= \lambda_2 [c_1, c_2] \stackrel{(6)}{=} 0. \end{aligned}$$

This is a contradiction.

Thus we have shown that $\mathcal{C}^1\mathfrak{g}$ cannot contain any J -invariant subspace which implies in turn that any complex structure will be abelian (Lemma 3.1 (viii)). $\square$

Lemma 3.12 — *Let $\mathfrak{g}$ be a 3-step nilpotent Lie-algebra with $\dim(\mathcal{C}^1\mathfrak{g}) = 3$ and $\dim(\mathcal{C}^2\mathfrak{g}) = \dim(\mathcal{C}^1\mathfrak{g} \cap \mathcal{Z}\mathfrak{g}) = 1$. Assume that J is a complex structure on $\mathfrak{g}$. Then $\mathcal{C}^2\mathfrak{g} \cap \mathcal{V}^J = 0$ but $\mathcal{V}^J \neq 0$ is possible and there may be nilpotent and non-nilpotent complex structures on the same Lie-algebra.*

Proof. Assume $\mathcal{C}^2\mathfrak{g} \cap \mathcal{V} \neq 0$. Then $(J\mathcal{Z}\mathfrak{g}) \cap \mathcal{C}^1\mathfrak{g} \neq 0$ because $\mathcal{V} \subset \mathcal{C}^1\mathfrak{g}$ and $\mathcal{C}^2\mathfrak{g} \subset \mathcal{Z}\mathfrak{g}$. Hence $\mathcal{V}_2 \neq 0$ which is impossible since $\dim \mathcal{C}^2\mathfrak{g} = 1$. We prove the second assertion in the following example. $\square$

Example 3.13 — Consider the following 10-dimensional Lie-algebra given by a basis $\mathfrak{g} = \langle e_1, e_2, e_3, e_4, e_5, e_6, z, c_1, c_2, c_3 \rangle$ and the structure equations

$$\begin{aligned} dc^1 &= e^{15} + e^{16} + e^{35} + e^{36} \\ dc^2 &= e^{25} + e^{26} + e^{45} + e^{46} \\ dc^3 &= e^1 \wedge c^1 + e^3 \wedge c^1 + e^2 \wedge c^2 + e^4 \wedge c^2 \end{aligned}$$

with respect to the dual basis. One can check that $d^2 = 0$ so the Jacobi-identity holds. The central filtrations are given by

$$\begin{aligned} \mathfrak{g} \supset \mathcal{Z}^2\mathfrak{g} &= \langle c_1, c_2, c_3, z \rangle \supset \mathcal{Z}^1\mathfrak{g} = \langle c_3, z \rangle \supset 0 \\ \mathfrak{g} \supset \mathcal{C}^1\mathfrak{g} &= \langle c_1, c_2, c_3 \rangle \supset \mathcal{C}^2\mathfrak{g} = \langle c_3 \rangle \supset 0 \end{aligned}$$

and hence we have $\dim \mathcal{C}^1 \mathfrak{g} = 3$, $\dim(\mathcal{Z} \mathfrak{g} \cap \mathcal{C}^1 \mathfrak{g}) = \dim \mathcal{C}^2 \mathfrak{g} = 1$. We give two complex structures J_1 and J_2 on $\mathfrak{g}$ by

$$\begin{aligned} J_1 e_1 &= e_2, & J_2 e_1 &= e_2, \\ J_1 e_3 &= e_4, & J_2 e_3 &= e_4, \\ J_1 e_5 &= e_6, & J_2 e_5 &= z, \\ J_1 c_1 &= c_2, & J_2 c_1 &= c_2, \\ J_1 z &= c_3, & J_2 e_6 &= c_3. \end{aligned}$$

A straightforward calculation shows that both structures are integrable and we have

$$\mathcal{V}^{J_1} = 0, \quad \mathcal{V}^{J_2} = \langle c_1, c_2 \rangle$$

which realises the two possibilities in Lemma 3.12. Note also that J_1 is nilpotent while J_2 is not nilpotent.

Lemma 3.14 — *Let $\mathfrak{g}$ be a 3-step nilpotent Lie-algebra with $\dim(\mathcal{C}^1 \mathfrak{g}) = 3$ and $\dim(\mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g}) = 2$.*

- (i) *If $\dim \mathcal{Z}^i \mathfrak{g}$ is even for all i , then every complex structure J on $\mathfrak{g}$ is nilpotent and one of the subspaces $\mathcal{Z} \mathfrak{g}$ and $\mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g}$ is J -invariant but there is no SPTBS in general.*
- (ii) *If $\dim \mathcal{Z}^1 \mathfrak{g}$ is odd or equal to 2 and $\dim \mathcal{Z}^2 \mathfrak{g}$ is even then*

$$\mathfrak{g} \supset \mathcal{Z}^2 \mathfrak{g} \supset \mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g} \supset 0$$

is a SPTBS for $\mathfrak{g}$.

- (iii) *If $\dim \mathcal{Z}^2 \mathfrak{g}$ is odd there does not exist any complex structure on $\mathfrak{g}$.*

Proof. Let $\mathfrak{g}$ be as above and assume that J is a complex structure on $\mathfrak{g}$. As a first step we show that $\mathcal{V} \subset \mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g}$.

Assume the contrary. For dimensional reasons we certainly have

$$\mathcal{V} \cap (\mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g}) \neq 0$$

so that we find some $z \in \mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g}$ such that $Jz \in \mathcal{C}^1 \mathfrak{g} \setminus (\mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g})$, i.e., $Jz \in \mathcal{W}_1 \setminus \mathcal{Z} \mathfrak{g}$ in the notation of Lemma 3.2 (ii). Thus $\mathcal{V}_2 \neq 0$ which is a contradiction if $\dim \mathcal{C}^2 \mathfrak{g} = 1$.

Else, if $\dim \mathcal{C}^2 \mathfrak{g} = 2$, both J -invariant subspaces coincide (Lemma 3.1 (iv)) and

$$\mathcal{V}_1 = \mathcal{V}_2 = \mathcal{C}^2 \mathfrak{g} = \mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g}$$

which contradicts our assumption $\mathcal{V} \not\subset \mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g}$.

Therefore we have $\mathcal{V} \subset \mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g}$ and at least one of the subspaces $\mathcal{Z} \mathfrak{g}$ and $\mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g}$ will be J -invariant. Note that the quotient of $\mathfrak{g}$ by any of these subspaces (which are ideals since they are contained in the centre) is a Lie-algebra with 1-dimensional commutator subalgebra. These have been studied in Section 3.2 and admit only abelian complex structures if the centre is evendimensional and no complex structures if it has odd dimension. This already proves (iii) and together with Example 3.15 also (i).

If $\dim \mathcal{Z} \mathfrak{g} = 2$ the two subspaces coincide and if $\dim \mathcal{Z} \mathfrak{g}$ is odd then $\mathcal{Z} \mathfrak{g}$ can never be invariant whence (ii). $\square$

In the following two examples we will show that all possibilities in (i) can indeed occur in one Lie-algebra and that we cannot achieve a better result by distinguishing the cases $\dim \mathcal{C}^2 \mathfrak{g} = 2$ and $\dim \mathcal{C}^2 \mathfrak{g} = 1$

Example 3.15 — Consider the following 8-dimensional 3-step nilpotent Lie-algebra given by a basis $\mathfrak{g} = \langle e_1, e_2, e_3, c_1, c_2, c_3, z_1, z_2 \rangle$ and the structure equations

$$\begin{aligned} dc^1 &= e^{12} \\ dc^2 &= e^1 \wedge c^1 + e^{23} \\ dc^3 &= e^2 \wedge c^1 - e^{13} \end{aligned}$$

with respect to the dual basis. This defines a Lie-algebra since $d^2 = 0$. The central filtrations are given by

$$\begin{aligned} \mathfrak{g} \supset \mathcal{Z}^2 \mathfrak{g} &= \langle e_3, c_1, c_2, c_3, z_1, z_2 \rangle \supset \mathcal{Z}^1 \mathfrak{g} = \langle c_2, c_3, z_1, z_2 \rangle \supset 0 \\ \mathfrak{g} \supset \mathcal{C}^1 \mathfrak{g} &= \langle c_1, c_2, c_3 \rangle \supset \mathcal{C}^2 \mathfrak{g} = \langle c_2, c_3 \rangle \supset 0 \end{aligned}$$

We give three complex structures J_1, J_2 and J_3 on $\mathfrak{g}$ by

$$\begin{array}{lll} J_1 e_1 = e_2, & J_2 e_1 = e_2, & J_3 e_1 = e_2, \\ J_1 c_1 = e_3, & J_2 c_1 = e_3, & J_3 c_2 = c_3, \\ J_1 c_3 = c_3, & J_2 c_2 = z_1, & J_3 z_1 = c_1, \\ J_1 z_1 = z_2, & J_2 c_3 = z_2, & J_3 z_2 = e_3. \end{array}$$

It is a straightforward calculation to check that these structures are integrable. The complex structure J_1 leaves both the commutator and the centre invariant, J_2 the centre but not the commutator, while J_3 leaves the commutator invariant but not the centre ($\mathcal{V}_{J_3} = \mathcal{C}^2 \mathfrak{g}$). This realises all possible combinations in Lemma 3.14 (i) in a single Lie-algebra.

Example 3.16 — Consider the following 18-dimensional Lie-algebra given by a basis $\mathfrak{g} = \langle e_1, e_2, e_3, f_1, f_2, f_3, g_1, g_2, g_3, h_1, h_2, h_3, z_1, z_2, c_0, \tilde{c}_0, c_1, c_2 \rangle$ and the structure equations

$$\begin{aligned} dc^0 &= g^{12} + g^{23} + h^{12} + h^{23} \\ dc^1 &= e^{12} + f^{12} \\ dc^2 &= e^{13} + f^{13} \\ &+ g^1 \wedge c^0 + c^0 \wedge g^3 + h^1 \wedge c^0 + c^0 \wedge h^3 \\ &- g^1 \wedge \tilde{c}^0 - \tilde{c}^0 \wedge g^3 + h^1 \wedge \tilde{c}^0 + \tilde{c}^0 \wedge h^3 \end{aligned}$$

with respect to the dual basis. The central filtrations are given by

$$\begin{aligned} \mathfrak{g} \supset \mathcal{Z}^2 \mathfrak{g} &= \langle e_1, e_2, e_3, f_1, f_2, f_3, c_0, \tilde{c}_0, c_1, c_2, z_1, z_2 \rangle \supset \mathcal{Z}^1 \mathfrak{g} = \langle c_1, c_2, z_1, z_2 \rangle \supset 0 \\ \mathfrak{g} \supset \mathcal{C}^1 \mathfrak{g} &= \langle c_0, c_1, c_2 \rangle \supset \mathcal{C}^2 \mathfrak{g} = \langle c_2 \rangle \supset 0 \end{aligned}$$

and we have $\dim(\mathcal{C}^1 \mathfrak{g}) = 3$, $\dim(\mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g}) = 2$ and $\dim(\mathcal{C}^2 \mathfrak{g}) = 1$.

We already studied the subalgebra

$$\mathfrak{a} = \langle e_1, e_2, e_3, f_1, f_2, f_3, z_1, z_2, c_1, c_2 \rangle$$

in Example 3.9 and in fact we have an extension

$$0 \rightarrow \mathfrak{a} \rightarrow \mathfrak{g} \rightarrow H_7 \oplus \mathbb{R} \rightarrow 0$$

where H_7 is the 7-dimensional Heisenberg algebra.

On the subspace $V = \langle g_1, g_2, g_3, h_1, h_2, h_3, c_0, \tilde{c}_0 \rangle$ complementary to $\mathfrak{a}$ we give a complex structure J by

$$Jc_0 = \tilde{c}_0, \quad Jg_i = h_i, \quad i = 1, \dots, 3.$$

Note that for $x, y \in V$ we have

$$[x, y] = [Jx, Jy] \text{ and } [x, Jy] = -[Jx, y]$$

and hence the Nijenhuis tensor $[x, y] - [Jx, Jy] + J([Jx, y] + [x, Jy])$ vanishes automatically for these elements in whatever way we choose to extend J to a complex structure on $\mathfrak{g}$. Furthermore we have $[V, \mathfrak{a}] = 0$ and therefore any integrable complex structure on $\mathfrak{a}$ can be combined with J to define an integrable complex structure on $\mathfrak{g}$.

Combining J with J_1, J_2 and J_3 from Example 3.9 we get three integrable complex structures on $\mathfrak{g}$ which realise the different possibilities in Lemma 3.14 (i).

4. APPLICATIONS

In this section we put together the results obtained so far. In particular we produce a number of classes of nilmanifolds which are closed under deformation in the large and completely describe the situation in real dimension six.

4.1. The Main Theorem. We want to apply Corollary 1.12 and Theorem 2.12 to our results obtained in Section 3. The outcome is the following:

Theorem 4.1 — *Let $M = (\mathfrak{g}, J, \Gamma)$ a nilmanifold with left-invariant complex structure. Then M and all small deformations of M are again iterated principal holomorphic torus bundles coming from a nilmanifold with left-invariant complex structure of type $(\mathfrak{g}, \Gamma)$ if one of the following conditions holds*

- $\dim \mathcal{C}^1 \mathfrak{g} \leq 2$.
- $\dim \mathcal{C}^1 \mathfrak{g} = 3$ and $\mathfrak{g}$ is 4-step nilpotent.
- $\dim \mathcal{C}^1 \mathfrak{g} = 3$, $\mathfrak{g}$ is 3-step nilpotent and $\dim(\mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z} \mathfrak{g}) = 2$.

Proof. We have shown in Proposition 3.6, 3.8 and 3.10 respectively that under the above conditions there is a principal torus bundle series for any given complex structure. Combining Theorem 1.10 with Theorem 1.11 we are done. $\square$

In order to proceed to deformations in the large, we need more good fibre classes:

Lemma 4.2 — *If $\mathfrak{h}$ is a nilpotent Lie-algebra with $\dim \mathcal{C}^1 \mathfrak{h} = 1$ then nilmanifolds with left-invariant complex structure of type $(\mathfrak{h}, \Gamma)$ constitute a good fibre class.*

Proof. Recall from Proposition 3.6 that all complex structures on $\mathfrak{g}$ are equivalent. Hence $h^1(M_J, \Theta) = h^{n-1,1}(M_J) = \dim H^{n-1,1}(\mathfrak{g}, J)$ is independent of the complex structure J . From the same proposition we know that $0 \subset \mathcal{Z}\mathfrak{h} \subset \mathfrak{h}$ is a SPTBS, and since complex tori are a good fibre class (see Remark 2.10), every deformation in the large is again of the same type by Theorem 2.12. Thus nilmanifolds with left-invariant complex structure of type $(\mathfrak{h}, \Gamma)$ satisfy both conditions for a good fibre class. $\square$

It would be desirable to have more good fibre classes but we can already prove

Theorem 4.3 — *Let $M = (\mathfrak{g}, J, \Gamma)$ a nilmanifold with left-invariant complex structure.*

(i) *M and all deformations in the large of M are iterated principal holomorphic torus bundles coming from a nilmanifold of type $(\mathfrak{g}, \Gamma)$ (with fibre dimensions depending only on $\mathfrak{g}$) if one of the following conditions holds:*

- $\mathfrak{g}$ is abelian.
- $\dim \mathcal{C}^1 \mathfrak{g} = 1$.
- $\mathfrak{g}$ satisfies the conditions of Proposition 3.4.
- $\dim \mathcal{C}^1 \mathfrak{g} = 2$, $\mathfrak{g}$ is 2-step nilpotent and $\dim(\mathcal{Z}\mathfrak{g})$ is equal to 2 or odd.
- $\dim \mathcal{C}^1 \mathfrak{g} = 2$ and $\mathfrak{g}$ is 3-step nilpotent.
- $\dim \mathcal{C}^1 \mathfrak{g} = 3$, $\mathfrak{g}$ is 3-step nilpotent and furthermore
 - $\dim(\mathcal{C}^1 \mathfrak{g} \cap \mathcal{Z}\mathfrak{g}) = 2$
 - $\dim \mathcal{Z}^1 \mathfrak{g}$ is odd or equal to 2.
 - $\dim \mathcal{Z}^2 \mathfrak{g}$ is even.
 - $\dim \mathcal{C}^1(\mathcal{Z}^2 \mathfrak{g}) = 1$.

The last condition is automatically fulfilled if $\dim \mathcal{C}^2 \mathfrak{g} = 1$.

(ii) *M and all small deformations of M are again iterated principal holomorphic torus bundles coming from a nilmanifold with left-invariant complex structure of type $(\mathfrak{g}, \Gamma)$ if one of the following conditions holds*

- $\mathfrak{g}$ admits a STBS $\mathfrak{g} \supset \mathcal{S}^1 \mathfrak{g} \supset 0$ of length 3.
- $\mathfrak{g}$ is 3-step nilpotent and admits a STBS of the form $\mathfrak{g} \supset \mathcal{S}^2 \mathfrak{g} \supset \mathcal{S}^1 \mathfrak{g} \supset 0$ such that $\dim(\mathcal{C}^1(\mathcal{S}^2 \mathfrak{g})) = 1$.

Proof. By Theorem 2.12, we have to check that in each of the cases there is a SPTBS (resp. STBS) $(\mathcal{S}^i \mathfrak{g})_{i=0, \dots, t}$ and that the nilmanifolds of the type $(\mathcal{S}^{t-1} \mathfrak{g}, \Gamma \cap \exp(\mathcal{S}^{t-1} \mathfrak{g}))$ constitute a good fibre class. The first assertion either holds by definition or has been proved in Proposition 3.4, 3.6, 3.8 and 3.10 respectively.

If $t = 1$ then M is a torus and the claim follows from [Cat04]. If $t = 2$, i.e., $\mathfrak{g}$ admits a SPTBS (or STBS)

$$\mathfrak{g} \supset \mathcal{S}^1 \mathfrak{g} \supset 0$$

then M is a holomorphic torus bundle over a torus and we are done, since tori constitute a good fibre class (see Remark 2.10). This is always the case if $\dim \mathcal{C}^1 \mathfrak{g} \leq 2$ and $\mathfrak{g}$ is not 3-step nilpotent and if $\mathfrak{g}$ satisfies the conditions of Proposition 3.4.

If $\mathfrak{g}$ is 3-step nilpotent and has a SPTBS $(\mathcal{S}^i \mathfrak{g})_{i=0,\dots,3}$, then we have to show that the nilmanifolds with Lie-algebra $\mathcal{S}^2 \mathfrak{g}$ constitute a good fibre class.

By Lemma 4.2 it suffices to check that $\dim \mathcal{C}^1(\mathcal{S}^2 \mathfrak{g}) = 1$ in the remaining cases. For a 3-step nilpotent Lie-algebra we have

$$\mathcal{C}^1(\mathcal{S}^2 \mathfrak{g}) \subset \mathcal{C}^1(\mathcal{Z}^2 \mathfrak{g}) \subset \mathcal{C}^2 \mathfrak{g}$$

and are done if $\dim \mathcal{C}^2 \mathfrak{g} = 1$. In the remaining cases the property holds by assumption. This concludes the proof. $\square$

4.2. Deformations and geometric structure in dimension 3. In this section we give a fairly complete classification of the geometric types of nilmanifolds with left-invariant complex structure in complex dimension 3 and determine their deformations.

In [Mag86] Magnin gave a classification of real nilpotent Lie-algebra in dimension at most 7 and in particular showed that in real dimension 6 there exist only 34 different isomorphism types while in higher dimension there are always continuous families.

Salamon showed in [Sal01] that only 18 of these 6-dimensional real nilpotent Lie-algebras admit a complex structure and Ugarte studied in detail the possible nilpotent and abelian structures ([Uga04], Theorem. 2.9); a part of these results has been reproved in Section 3. Following Ugarte's notation we give the list of 6-dimensional real Lie-algebras admitting complex structures:

$$\begin{aligned} \mathfrak{h}_1 &= (0, 0, 0, 0, 0, 0), & \mathfrak{h}_{10} &= (0, 0, 0, 12, 13, 14), \\ \mathfrak{h}_2 &= (0, 0, 0, 0, 12, 34), & \mathfrak{h}_{11} &= (0, 0, 0, 12, 13, 14 + 23), \\ \mathfrak{h}_3 &= (0, 0, 0, 0, 0, 12 + 34), & \mathfrak{h}_{12} &= (0, 0, 0, 12, 13, 24), \\ \mathfrak{h}_4 &= (0, 0, 0, 0, 12, 14 + 23), & \mathfrak{h}_{13} &= (0, 0, 0, 12, 13 + 14, 24), \\ \mathfrak{h}_5 &= (0, 0, 0, 0, 13 + 42, 14 + 23), & \mathfrak{h}_{14} &= (0, 0, 0, 12, 14, 13 + 42), \\ \mathfrak{h}_6 &= (0, 0, 0, 0, 12, 13), & \mathfrak{h}_{15} &= (0, 0, 0, 12, 13 + 42, 14 + 23), \\ \mathfrak{h}_7 &= (0, 0, 0, 12, 13, 23), & \mathfrak{h}_{16} &= (0, 0, 0, 12, 14, 24), \\ \mathfrak{h}_8 &= (0, 0, 0, 0, 0, 12), & \mathfrak{h}_{19}^- &= (0, 0, 0, 12, 23, 14 - 35), \\ \mathfrak{h}_9 &= (0, 0, 0, 0, 12, 14 + 25), & \mathfrak{h}_{26}^+ &= (0, 0, 12, 13, 23, 14 + 25). \end{aligned}$$

Our classification then reads as follows:

Theorem 4.4 — *Let $M_J = (\mathfrak{g}, J, \Gamma)$ be a complex 3-dimensional nilmanifold with left-invariant complex structure.*

- (i) *If $\mathfrak{g}$ is not in $\{\mathfrak{h}_7, \mathfrak{h}_{19}^-, \mathfrak{h}_{26}^+\}$, then $\mathfrak{g}$ admits a SPTBS and hence M_J has the structure of an iterated principal holomorphic torus bundle. We list the possibilities in the following table:*

base	fibre	corresponding Lie-algebras
3-torus	-	$\mathfrak{h}_1$
2-torus	elliptic curve	$\mathfrak{h}_2, \mathfrak{h}_3, \mathfrak{h}_4, \mathfrak{h}_5, \mathfrak{h}_6$
elliptic curve	2-torus	$\mathfrak{h}_8$
Kodaira surface	elliptic curve	$\mathfrak{h}_9, \mathfrak{h}_{10}, \mathfrak{h}_{11}, \mathfrak{h}_{12}, \mathfrak{h}_{13}, \mathfrak{h}_{14}, \mathfrak{h}_{15}, \mathfrak{h}_{16}$

Every deformation in the large is of the same type.

- (ii) If $\mathfrak{g} = \mathfrak{h}_{19}^-$ or $\mathfrak{g} = \mathfrak{h}_{26}^+$ then $\mathfrak{g}$ admits a STBS and M_J can be described as 2-torus bundle over an elliptic curve but there is no principal torus bundle structure. Every deformation in the large is of the same type.
- (iii) If $\mathfrak{g} = \mathfrak{h}_7$ then there is a dense subset of the space of all left-invariant complex structures for which M admits the structure of principal holomorphic bundle of elliptic curves over a Kodaira surface but this is not true for all complex structures.

Proof. First of all note that every nilpotent Lie-algebra of real dimension at most 4 which admits complex structures gives rise to a good fibre class, since the only possibilities are elliptic curves, 2-dimensional tori and Kodaira surfaces. So for (i) and (ii) it suffices to exhibit SPTBS resp. STBS with the appropriate dimensions.

We start with the cases in (i). The calculation of the dimensions of the descending and ascending series is straightforward and we find that all cases are covered by the results in Section 3:

Proposition	3.2	3.8 (i)	3.8 (ii)	3.8 (iii)	3.10 (iii) (b)
Lie-algebra	$\mathfrak{h}_3, \mathfrak{h}_8$	$\mathfrak{h}_9$	$\mathfrak{h}_6$	$\mathfrak{h}_2, \mathfrak{h}_4, \mathfrak{h}_5$	$\mathfrak{h}_{10}, \mathfrak{h}_{11}, \mathfrak{h}_{12}, \mathfrak{h}_{13},$ $\mathfrak{h}_{14}, \mathfrak{h}_{15}, \mathfrak{h}_{16}$

The Lie-algebra $\mathfrak{h}_1$ corresponds to the complex 3-dimensional torus.

The Lie-algebras $\mathfrak{h}_{19}^-$ and $\mathfrak{h}_{26}^+$ do not admit any nilpotent complex structure (see [Sal01] or note that the centre has real dimension 1) and therefore a corresponding nilmanifold can never admit an iterated principal holomorphic torus bundle structure. Thus it remains to exhibit a STBS in both cases.

The case $\mathfrak{h}_{26}^+$: We claim that $\mathfrak{g} \supset \mathcal{C}^1 \mathfrak{g} := \langle e_3, \dots, e_6 \rangle \supset 0$ is a stable torus bundle series. Since the commutator is always a rational subspace with respect to every rational structure, we only need prove that for every complex structure J on $\mathfrak{h}_{26}^+$ the subspace $\mathcal{C}^1 \mathfrak{g}$ is J -invariant.

Let $Je_6 = \sum i = 1^6 \lambda_i e_i$, $W := \mathcal{Z}_J \mathfrak{h}_{26}^+ = \langle e_6, Je_6 \rangle$ and, as in Lemma 3.2, $\mathcal{V} := \text{im } ad_{Je_6}$. By Lemma 3.5 we know that $\mathcal{V} \cap W = 0$ and, applying ad_{Je_6} to e_4 and e_5 , we see that $\lambda_1 = \lambda_2 = 0$. As $W \subset \mathcal{C}^1 \mathfrak{g}$, $\mathcal{V} \subset \mathcal{C}^1 \mathfrak{g}$ and their intersection is 0 so $\mathcal{C}^1 \mathfrak{g} = \mathcal{V} \oplus W$ is J -invariant.

The case $\mathfrak{h}_{19}^-$: We claim that $\mathfrak{g} \supset V := \langle e_2, e_4, e_5, e_6 \rangle \supset 0$ is a stable torus bundle series. With the same proof as in the first case (replacing e_2 by e_3) we see that V is J -invariant for every complex structure in $\mathfrak{h}_{19}^-$.

It is straightforward to check that

$$V = \{x \in \mathfrak{h}_{19}^- \mid (ad_x)^2 = 0\}$$

which implies that V is rational with respect to every rational structure.

It remains to prove (iii). A nilmanifold of type $M_J = (\mathfrak{h}_7, J, \Gamma)$ admits a structure of iterated principal holomorphic torus bundle if and only if the subspaces $\mathcal{T}^1 \mathfrak{h}_7$ and $\mathcal{T}^2 \mathfrak{h}_7$ are Γ -rational. This is clearly the case if the complex structure is rational but not always as seen in Example 1.14. A

quotient of $\mathfrak{h}_7$ by a 2-dimensional subspace of the centre can not be abelian, so $\mathcal{T}^2\mathfrak{h}_7 \neq \mathfrak{h}_7$ and, if there is any iterated bundle structure, we have a bundle of elliptic curves over a Kodaira surface. $\square$

Remark 4.5 — The real Lie-algebra underlying the Iwasawa manifold is isomorphic to $\mathfrak{h}_5$ and hence we have in particular proved that every deformation in the large of the Iwasawa manifold is a nilmanifold with left-invariant complex structure. It would be interesting to know what are the deformations in the large of nilmanifolds with Lie-algebra $\mathfrak{h}_7$.

REFERENCES

- [BG88] Chal Benson and Carolyn S. Gordon. Kähler and symplectic structures on nilmanifolds. *Topology*, 27(4):513–518, 1988.
- [Bor84] Ciprian Borcea. Moduli for Kodaira surfaces. *Compositio Math.*, 52(3):373–380, 1984.
- [Cat91] Fabrizio Catanese. Moduli and classification of irregular Kaehler manifolds (and algebraic varieties) with Albanese general type fibrations. *Invent. Math.*, 104(2):263–289, 1991.
- [Cat02] Fabrizio Catanese. Deformation types of real and complex manifolds. In *Contemporary trends in algebraic geometry and algebraic topology (Tianjin, 2000)*, volume 5 of *Nankai Tracts Math.*, pages 195–238. World Sci. Publ., River Edge, NJ, 2002.
- [Cat04] Fabrizio Catanese. Deformation in the large of some complex manifolds. I. *Ann. Mat. Pura Appl. (4)*, 183(3):261–289, 2004.
- [Cat07] Fabrizio Catanese. Q.E.D. for algebraic varieties. *J. Differential Geom.*, 77(1):43–75, 2007.
- [CF01] S. Console and A. Fino. Dolbeault cohomology of compact nilmanifolds. *Transform. Groups*, 6(2):111–124, 2001.
- [CF06] Fabrizio Catanese and Paola Frediani. Deformation in the large of some complex manifolds. II. In *Recent progress on some problems in several complex variables and partial differential equations*, volume 400 of *Contemp. Math.*, pages 21–41. Amer. Math. Soc., Providence, RI, 2006.
- [CFGU00] Luis A. Cordero, Marisa Fernández, Alfred Gray, and Luis Ugarte. Compact nilmanifolds with nilpotent complex structures: Dolbeault cohomology. *Trans. Amer. Math. Soc.*, 352(12):5405–5433, 2000.
- [CFP06] S. Console, A. Fino, and Y. S. Poon. Stability of abelian complex structures. *Internat. J. Math.*, 17(4):401–416, 2006.
- [CG90] Lawrence J. Corwin and Frederick P. Greenleaf. *Representations of nilpotent Lie groups and their applications. Part I*, volume 18 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1990. Basic theory and examples.
- [FG65] Wolfgang Fischer and Hans Grauert. Lokal-triviale Familien kompakter komplexer Mannigfaltigkeiten. *Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. II*, 1965:89–94, 1965.
- [GMPP04] Gueo Grantcharov, Colin McLaughlin, Henrik Pedersen, and Yat Sun Poon. Deformations of Kodaira manifolds. *Glasg. Math. J.*, 46(2):259–281, 2004.
- [GR84] Hans Grauert and Reinhold Remmert. *Coherent analytic sheaves*, volume 265 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1984.
- [Has89] Keizo Hasegawa. Minimal models of nilmanifolds. *Proc. Amer. Math. Soc.*, 106(1):65–71, 1989.
- [Huy05] Daniel Huybrechts. *Complex geometry*. Universitext. Springer-Verlag, Berlin, 2005.
- [KN69] Shoshichi Kobayashi and Katsumi Nomizu. *Foundations of differential geometry. Vol. II*. Interscience Tracts in Pure and Applied Mathematics, No. 15 Vol.

- II. Interscience Publishers John Wiley & Sons, Inc., New York-London-Sydney, 1969.
- [KS58] K. Kodaira and D. C. Spencer. On deformations of complex analytic structures. I, II. *Ann. of Math. (2)*, 67:328–466, 1958.
 - [KS04] Georgios Katsenzis and Simon Salamon. Complex structures on the Iwasawa manifold. *Adv. Geom.*, 4(2):165–179, 2004.
 - [Kur62] M. Kuranishi. On the locally complete families of complex analytic structures. *Ann. of Math. (2)*, 75:536–577, 1962.
 - [Lan84] Serge Lang. *Algebra*. Addison-Wesley Publishing Company Advanced Book Program, Reading, MA, second edition, 1984.
 - [Mag86] L. Magnin. Sur les algèbres de Lie nilpotentes de dimension ≤ 7 . *J. Geom. Phys.*, 3(1):119–144, 1986.
 - [Mal51] A. I. Malcev. On a class of homogeneous spaces. *Amer. Math. Soc. Translation*, 1951(39):33, 1951.
 - [MPPS06] C. Maclaughlin, H. Pedersen, Y. S. Poon, and S. Salamon. Deformation of 2-step nilmanifolds with abelian complex structures. *J. London Math. Soc. (2)*, 73(1):173–193, 2006.
 - [Nom54] Katsumi Nomizu. On the cohomology of compact homogeneous spaces of nilpotent Lie groups. *Ann. of Math. (2)*, 59:531–538, 1954.
 - [Rol07a] Sönke Rollenske. *Nilmanifolds: Complex structures, geometry and deformations*. PhD thesis, Universität Bayreuth, 2007.
 - [Rol07b] Sönke Rollenske. Q.E.D. for Kodaira surfaces, 2007. Appendix to [Cat07].
 - [Rol07c] Sönke Rollenske. The Frölicher spectral sequence can be arbitrarily non degenerate, 2007. to appear in *Mathematische Annalen*.
 - [Rol08a] Sönke Rollenske. The Kuranishi space of complex parallelisable nilmanifolds, 2008, arXiv:0803.2048v1.
 - [Rol08b] Sönke Rollenske. Lie-algebra Dolbeault cohomology and small deformations of nilmanifolds, 2008. preprint.
 - [Sak76] Yusuke Sakane. On compact complex parallelisable solvmanifolds. *Osaka J. Math.*, 13(1):187–212, 1976.
 - [Sal01] S. M. Salamon. Complex structures on nilpotent Lie algebras. *J. Pure Appl. Algebra*, 157(2-3):311–333, 2001.
 - [Uga04] Luis Ugarte. Hermitian structures on six dimensional nilmanifolds, 2004, arXiv:math.DG/0411254.
 - [Var84] V. S. Varadarajan. *Lie groups, Lie algebras, and their representations*, volume 102 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1984. Reprint of the 1974 edition.
 - [VGS00] E. B. Vinberg, V. V. Gorbatsevich, and O. V. Shvartsman. Discrete subgroups of Lie groups [MR0968445 (90c:22036)]. In *Lie groups and Lie algebras, II*, volume 21 of *Encyclopaedia Math. Sci.*, pages 1–123, 217–223. Springer, Berlin, 2000.
 - [Wav69] John J. Wavrik. Obstructions to the existence of a space of moduli. In *Global Analysis (Papers in Honor of K. Kodaira)*, pages 403–414. Univ. Tokyo Press, Tokyo, 1969.
 - [Win98] Jörg Winkelmann. Complex analytic geometry of complex parallelizable manifolds. *Mém. Soc. Math. Fr. (N.S.)*, (72-73):x+219, 1998.

DR. SÖNKE ROLLENSKE, DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, SW7 2AZ LONDON, UNITED KINGDOM

E-mail address: s.rollenske@imperial.ac.uk